



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>5</sup> :  <b>C12N 15/12, C07K 13/00</b>  <b>C12N 5/10, A61K 37/02</b>  <b>G01N 33/68</b></p>	<b>A2</b>	<p>(11) International Publication Number: <b>WO 94/11502</b></p> <p>(43) International Publication Date: <b>26 May 1994 (26.05.94)</b></p>																																																																															
<p>(21) International Application Number: <b>PCT/GB93/02367</b></p> <p>(22) International Filing Date: <b>17 November 1993 (17.11.93)</b></p> <p>(30) Priority data:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">9224057.1</td> <td style="width: 30%;">17 November 1992 (17.11.92)</td> <td style="width: 40%;">GB</td> </tr> <tr> <td>9304677.9</td> <td>8 March 1993 (08.03.93)</td> <td>GB</td> </tr> <tr> <td>9304680.3</td> <td>8 March 1993 (08.03.93)</td> <td>GB</td> </tr> <tr> <td>9311047.6</td> <td>28 May 1993 (28.05.93)</td> <td>GB</td> </tr> <tr> <td>9313763.6</td> <td>2 July 1993 (02.07.93)</td> <td>GB</td> </tr> <tr> <td>9316099.2</td> <td>3 August 1993 (03.08.93)</td> <td>GB</td> </tr> <tr> <td>9321344.5</td> <td>15 October 1993 (15.10.93)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except US): <b>LUDWIG INSTITUTE FOR CANCER RESEARCH [GB/GB]; St. Mary's Hospital Medical School, Norfolk Place, Paddington, London W2 1PG (GB).</b></p>			9224057.1	17 November 1992 (17.11.92)	GB	9304677.9	8 March 1993 (08.03.93)	GB	9304680.3	8 March 1993 (08.03.93)	GB	9311047.6	28 May 1993 (28.05.93)	GB	9313763.6	2 July 1993 (02.07.93)	GB	9316099.2	3 August 1993 (03.08.93)	GB	9321344.5	15 October 1993 (15.10.93)	GB																																																										
9224057.1	17 November 1992 (17.11.92)	GB																																																																															
9304677.9	8 March 1993 (08.03.93)	GB																																																																															
9304680.3	8 March 1993 (08.03.93)	GB																																																																															
9311047.6	28 May 1993 (28.05.93)	GB																																																																															
9313763.6	2 July 1993 (02.07.93)	GB																																																																															
9316099.2	3 August 1993 (03.08.93)	GB																																																																															
9321344.5	15 October 1993 (15.10.93)	GB																																																																															
<p>(72) Inventors; and          (75) Inventors/Applicants (for US only) : <b>MIYAZONO, Kohei [JP/SE]; Flogstavägen 63D, S-752 63 Uppsala (SE). DIJKE, Peter, Ten [NL/SE]; Flogstavägen 25C, S-752 63 Uppsala (SE). FRANZEN, Petra [SE/SE]; Lindsbergsgatan 15b, S-752 40 Uppsala (SE). YAMASHITA, Hidetoshi [JP/SE]; Flogstavägen 33A, S-752 63 Uppsala (SE). HELDIN, Carl-Henrik [SE/SE]; Hesselmanns väg 35, S-752 63 Uppsala (SE).</b></p> <p>(74) Agent: <b>GILL JENNINGS &amp; EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).</b></p> <p>(81) Designated States: <b>AU, CA, JP, KR, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b></p> <p><b>Published</b>  <i>Without international search report and to be republished upon receipt of that report.</i></p> <p style="text-align: center; font-weight: bold; font-size: 1.2em;">BEST AVAILABLE COPY</p>																																																																																	
<p>(54) Title: <b>ACTIVIN RECEPTOR-LIKE KINASES, PROTEINS HAVING SERINE THREONINE KINASE DOMAINS AND THEIR USE</b></p>																																																																																	
<table style="width: 100%; border: none;"> <tr> <td style="width: 20%;">cons.aa</td> <td style="width: 20%; text-align: center;">G G G V</td> <td style="width: 20%; text-align: center;">A K</td> <td style="width: 20%; text-align: center;">E</td> <td style="width: 40%;"></td> </tr> <tr> <td>htGFBR-II</td> <td>LDTLVGKGRFAEVYKAKLKQNTSEQFETVAVKIFPYDHYASWDRKDI</td> <td></td> <td></td> <td>IFSDINLKHENILQF</td> </tr> <tr> <td>mActR-IIB</td> <td>LLEIKARGRFPCVWKAQLMN-----</td> <td>DFVAVKIKPLQDKQSWOSEREI</td> <td></td> <td>FSTPGMGHENLLQF</td> </tr> <tr> <td>mActR-II</td> <td>LLEVKARGRFPCVWKAQLLN-----</td> <td>EYVAVKIFPIQDKQSWQNEYEVYSI</td> <td></td> <td>PGMGHENILQF</td> </tr> <tr> <td>daf-1</td> <td>LQVRVSGSRFGNVSRGDYRG-----</td> <td>EAVAVKVFNADPAFHKEIEI</td> <td></td> <td>PETRMRLRHPNVLR</td> </tr> <tr> <td>subdomains</td> <td style="text-align: center;">I</td> <td style="text-align: center;">II</td> <td style="text-align: center;">III</td> <td style="text-align: center;">IV</td> </tr> </table> <table style="width: 100%; border: none;"> <tr> <td style="width: 20%;">htGFBR-II</td> <td style="width: 20%;">LTAERKTELKQYWLITAFHAKGNLQEYLTRHVISWEDLRNVGSSLARGLSHLSDHTP-C</td> <td style="width: 20%;"></td> <td style="width: 20%;"></td> <td style="width: 40%;"></td> </tr> <tr> <td>mActR-IIB</td> <td>IAAEKRGSNLEVELMLITAFHDKGSLIDYLGKNIITWNECHVAETMSRGISYLHEDVPWCR</td> <td></td> <td></td> <td></td> </tr> <tr> <td>mActR-II</td> <td>IGAERKGTSDVDLMLITAFHEKGSLSDFLEKANNVSWNELCHIAETMARGLAYLHEDI</td> <td></td> <td></td> <td>PGGLK</td> </tr> <tr> <td>daf-1</td> <td>IGSDRVDTGFTVTEMLVIEYHPSGSLHDFLENTVNIETTYNLMRSTASGLAFLNQIGGSK</td> <td></td> <td></td> <td></td> </tr> <tr> <td>subdomains</td> <td style="text-align: center;">V</td> <td style="text-align: center;">VI-A</td> <td colspan="2"></td> </tr> </table> <table style="width: 100%; border: none;"> <tr> <td style="width: 20%;">cons.aa</td> <td style="width: 20%; text-align: center;">DLK N</td> <td style="width: 20%; text-align: center;">DFG</td> <td style="width: 40%;"></td> </tr> <tr> <td>htGFBR-II</td> <td>-GRPMPIVHRDLKSSNLLVKNLITCCLCDFGLSLRL--</td> <td>GPYSSVDDLANSQGVGTARYMAP</td> <td></td> </tr> <tr> <td>mActR-IIB</td> <td>GEGHKPSIAHRDFKSNVLLKSDLTAVLADFGLAVERF--</td> <td>EPGKPPGD--</td> <td>THQGVGTRRYMAP</td> </tr> <tr> <td>mActR-II</td> <td>-DGHKPAISHRDIKSNVLLKNNLTACTADFGALALF--</td> <td>EAGKSAGD--</td> <td>THQGVGTRRYMAP</td> </tr> <tr> <td>daf-1</td> <td>-ESNKPAMAHARDIKSNIMYKNDLTCAIGDLGLSLSKPEDAASDI</td> <td>IAN--</td> <td>ENYKCGTVRYLAP</td> </tr> <tr> <td>subdomains</td> <td style="text-align: center;">VI-B</td> <td style="text-align: center;">VII</td> <td style="text-align: center;">VIII</td> </tr> </table>			cons.aa	G G G V	A K	E		htGFBR-II	LDTLVGKGRFAEVYKAKLKQNTSEQFETVAVKIFPYDHYASWDRKDI			IFSDINLKHENILQF	mActR-IIB	LLEIKARGRFPCVWKAQLMN-----	DFVAVKIKPLQDKQSWOSEREI		FSTPGMGHENLLQF	mActR-II	LLEVKARGRFPCVWKAQLLN-----	EYVAVKIFPIQDKQSWQNEYEVYSI		PGMGHENILQF	daf-1	LQVRVSGSRFGNVSRGDYRG-----	EAVAVKVFNADPAFHKEIEI		PETRMRLRHPNVLR	subdomains	I	II	III	IV	htGFBR-II	LTAERKTELKQYWLITAFHAKGNLQEYLTRHVISWEDLRNVGSSLARGLSHLSDHTP-C				mActR-IIB	IAAEKRGSNLEVELMLITAFHDKGSLIDYLGKNIITWNECHVAETMSRGISYLHEDVPWCR				mActR-II	IGAERKGTSDVDLMLITAFHEKGSLSDFLEKANNVSWNELCHIAETMARGLAYLHEDI			PGGLK	daf-1	IGSDRVDTGFTVTEMLVIEYHPSGSLHDFLENTVNIETTYNLMRSTASGLAFLNQIGGSK				subdomains	V	VI-A			cons.aa	DLK N	DFG		htGFBR-II	-GRPMPIVHRDLKSSNLLVKNLITCCLCDFGLSLRL--	GPYSSVDDLANSQGVGTARYMAP		mActR-IIB	GEGHKPSIAHRDFKSNVLLKSDLTAVLADFGLAVERF--	EPGKPPGD--	THQGVGTRRYMAP	mActR-II	-DGHKPAISHRDIKSNVLLKNNLTACTADFGALALF--	EAGKSAGD--	THQGVGTRRYMAP	daf-1	-ESNKPAMAHARDIKSNIMYKNDLTCAIGDLGLSLSKPEDAASDI	IAN--	ENYKCGTVRYLAP	subdomains	VI-B	VII	VIII
cons.aa	G G G V	A K	E																																																																														
htGFBR-II	LDTLVGKGRFAEVYKAKLKQNTSEQFETVAVKIFPYDHYASWDRKDI			IFSDINLKHENILQF																																																																													
mActR-IIB	LLEIKARGRFPCVWKAQLMN-----	DFVAVKIKPLQDKQSWOSEREI		FSTPGMGHENLLQF																																																																													
mActR-II	LLEVKARGRFPCVWKAQLLN-----	EYVAVKIFPIQDKQSWQNEYEVYSI		PGMGHENILQF																																																																													
daf-1	LQVRVSGSRFGNVSRGDYRG-----	EAVAVKVFNADPAFHKEIEI		PETRMRLRHPNVLR																																																																													
subdomains	I	II	III	IV																																																																													
htGFBR-II	LTAERKTELKQYWLITAFHAKGNLQEYLTRHVISWEDLRNVGSSLARGLSHLSDHTP-C																																																																																
mActR-IIB	IAAEKRGSNLEVELMLITAFHDKGSLIDYLGKNIITWNECHVAETMSRGISYLHEDVPWCR																																																																																
mActR-II	IGAERKGTSDVDLMLITAFHEKGSLSDFLEKANNVSWNELCHIAETMARGLAYLHEDI			PGGLK																																																																													
daf-1	IGSDRVDTGFTVTEMLVIEYHPSGSLHDFLENTVNIETTYNLMRSTASGLAFLNQIGGSK																																																																																
subdomains	V	VI-A																																																																															
cons.aa	DLK N	DFG																																																																															
htGFBR-II	-GRPMPIVHRDLKSSNLLVKNLITCCLCDFGLSLRL--	GPYSSVDDLANSQGVGTARYMAP																																																																															
mActR-IIB	GEGHKPSIAHRDFKSNVLLKSDLTAVLADFGLAVERF--	EPGKPPGD--	THQGVGTRRYMAP																																																																														
mActR-II	-DGHKPAISHRDIKSNVLLKNNLTACTADFGALALF--	EAGKSAGD--	THQGVGTRRYMAP																																																																														
daf-1	-ESNKPAMAHARDIKSNIMYKNDLTCAIGDLGLSLSKPEDAASDI	IAN--	ENYKCGTVRYLAP																																																																														
subdomains	VI-B	VII	VIII																																																																														
<p>(57) Abstract</p> <p>A new receptor family has been identified, of activin-like kinases. Novel proteins have activin/TGF-<math>\beta</math>-type I receptor functionality, and have consequential diagnostic/therapeutic utility. They may have a serine/threonine kinase domain, a DFKSRN or DLKSKN sequence in subdomain VIB and/or a GTKRYM sequence in subdomain VIII.</p>																																																																																	

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ACTIVIN RECEPTOR-LIKE KINASES, PROTEINS HAVING  
SERINE THREONINE KINASE DOMAINS AND THEIR USE.

\*\*\*\*\*

Field of the Invention

5 This invention relates to proteins having serine/threonine kinase domains, corresponding nucleic acid molecules, and their use.

Background of the Invention

10 The transforming growth factor- $\beta$  (TGF- $\beta$ ) superfamily consists of a family of structurally-related proteins, including three different mammalian isoforms of TGF- $\beta$  (TGF- $\beta$ 1,  $\beta$ 2 and  $\beta$ 3), activins, inhibins, müllerian-inhibiting substance and bone morphogenic proteins (BMPs) (for reviews see Roberts and Sporn, (1990) Peptide Growth Factors and Their Receptors, Pt.1, Sporn and Roberts, eds. (Berlin: Springer - Verlag) pp 419-472; Moses et al (1990) Cell 63, 245-247). The proteins of the TGF- $\beta$  superfamily have a wide variety of biological activities. TGF- $\beta$  acts as a growth inhibitor for many cell types and appears to play a central role in the regulation of embryonic development, 20 tissue regeneration, immuno-regulation, as well as in fibrosis and carcinogenesis (Roberts and Sporn (199) see above).

25 Activins and inhibins were originally identified as factors which regulate secretion of follicle-stimulating hormone secretion (Vale et al (1990) Peptide Growth Factors and Their Receptors, Pt.2, Sporn and Roberts, eds. (Berlin: Springer-Verlag) pp.211-248). Activins were also shown to induce the differentiation of haematopoietic progenitor cells (Murata et al (1988) Proc. Natl. Acad. Sci. USA 85, 2434 - 2438; Eto et al (1987) Biochem. Biophys. Res. Commun. 142, 1095-1103) and induce mesoderm formation in Xenopus embryos (Smith et al (1990) Nature 345, 729-731; van den Eijnden-Van Raaij et al (1990) Nature 345, 732-734).

35 BMPs or osteogenic proteins which induce the formation of bone and cartilage when implanted subcutaneously (Wozney et al (1988) Science 242, 1528-1534), facilitate neuronal

differentiation (Paralkar et al (1992) J. Cell Biol. 119, 1721-1728) and induce monocyte chemotaxis (Cunningham et al (1992) Proc. Natl. Acad. Sci. USA 89, 11740-11744). Müllerian-inhibiting substance induces regression of the  
5 Müllerian duct in the male reproductive system (Cate et al (1986) Cell 45, 685-698), and a glial cell line-derived neurotrophic factor enhances survival of midbrain dopaminergic neurons (Lin et al (1993) Science 260, 1130-1132). The action of these growth factors is mediated  
10 through binding to specific cell surface receptors.

Within this family, TGF- $\beta$  receptors have been most thoroughly characterized. By covalently cross-linking radio-labelled TGF- $\beta$  to cell surface molecules followed by polyacrylamide gel electrophoresis of the affinity-labelled  
15 complexes, three distinct size classes of cell surface proteins (in most cases) have been identified, denoted receptor type I (53 kd), type II (75 kd), type III or betaglycan (a 300 kd proteoglycan with a 120 kd core protein) (for a review see Massague (1992) Cell 69 1067-1070) and more recently endoglin (a homodimer of two 95 kd subunits) (Cheifetz et al (1992) J. Biol. Chem. 267 19027-19030). Current evidence suggests that type I and type II receptors are directly involved in receptor signal transduction (Segarini et al (1989) Mol. Endo., 3, 261-272; Laiho et al (1991) J. Biol. Chem. 266, 9100-9112) and may  
25 form a heteromeric complex; the type II receptor is needed for the binding of TGF- $\beta$  to the type I receptor and the type I receptor is needed for the signal transduction induced by the type II receptor (Wrana et al (1992) Cell, 71, 1003-1004). The type III receptor and endoglin may have more indirect roles, possibly by facilitating the binding of ligand to type II receptors (Wang et al (1991) Cell, 67 797-805; López-Casillas et al (1993) Cell, 73 1435-1444).

35 Binding analyses with activin A and BMP4 have led to the identification of two co-existing cross-linked affinity complexes of 50-60 kDa and 70-80 kDa on responsive cells

(Hino et al (1989) J. Biol. Chem. 264, 10309 - 10314; Mathews and Vale (1991), Cell 68, 775-785; Paralker et al (1991) Proc. Natl. Acad. Sci. USA 87, 8913-8917). By analogy with TGF- $\beta$  receptors they are thought to be signalling receptors and have been named type I and type II receptors.

Among the type II receptors for the TGF- $\beta$  superfamily of proteins, the cDNA for the activin type II receptor (ActRII) was the first to be cloned (Mathews and Vale (1991) Cell 65, 973-982). The predicted structure of the receptor was shown to be a transmembrane protein with an intracellular serine/threonine kinase domain. The activin receptor is related to the C. elegans daf-1 gene product, but the ligand is currently unknown (Georgi et al (1990) Cell 61, 635-645). Thereafter, another form of the activin type II receptor (activin type IIB receptor), of which there are different splicing variants (Mathews et al (1992), Science 225, 1702-1705; Attisano et al (1992) Cell 68, 97-108), and the TGF- $\beta$  type II receptor (TBRII) (Lin et al (1992) Cell 68, 775-785) were cloned, both of which have putative serine/threonine kinase domains.

#### Summary of the Invention

The present invention involves the discovery of related novel peptides, including peptides having the activity of those defined herein as SEQ ID Nos. 2, 4, 8, 10, 12, 14, 16 and 18. Their discovery is based on the realisation that receptor serine/threonine kinases form a new receptor family, which may include the type II receptors for other proteins in the TGF- $\beta$  superfamily. To ascertain whether there were other members of this family of receptors, a protocol was designed to clone ActRII/daf I related cDNAs. This approach made use of the polymerase chain reaction (PCR), using degenerate primers based upon the amino-acid sequence similarity between kinase domains of the mouse activin type II receptor and daf-I gene products.

This strategy resulted in the isolation of a new family of receptor kinases called Activin receptor like kinases (ALK's) 1-6. These cDNAs showed an overall 33-39% sequence similarity with ActRII and TGF- $\beta$  type II receptor and 40-92% sequence similarity towards each other in the kinase domains.

Soluble receptors according to the invention comprise at least predominantly the extracellular domain. These can be selected from the information provided herein, prepared in conventional manner, and used in any manner associated with the invention.

Antibodies to the peptides described herein may be raised in conventional manner. By selecting unique sequences of the peptides, antibodies having desired specificity can be obtained.

The antibodies may be monoclonal, prepared in known manner. In particular, monoclonal antibodies to the extracellular domain are of potential value in therapy.

Products of the invention are useful in diagnostic methods, e.g. to determine the presence in a sample for an analyte binding therewith, such as in an antagonist assay. Conventional techniques, e.g. an enzyme-linked immunosorbent assay, may be used.

Products of the invention having a specific receptor activity can be used in therapy, e.g. to modulate conditions associated with activin or TGF- $\beta$  activity. Such conditions include fibrosis, e.g. liver cirrhosis and pulmonary fibrosis, cancer, rheumatoid arthritis and glomeronephritis.

### Brief Description of the Drawings

Figure 1 shows the alignment of the serine/threonine (S/T) kinase domains (I-VIII) of related receptors from transmembrane proteins, including embodiments of the present invention. The nomenclature of the subdomains is accordingly to Hanks et al (1988).

Figures 2A to 2D shows the sequences and characteristics of the respective primers used in the

initial PCR reactions. The nucleic acid sequences are also given as SEQ ID Nos. 19 to 22.

Figure 3 is a comparison of the amino-acid sequences of human activin type II receptor (Act R-II), mouse activin type IIB receptor (Act R-IIB), human TGF- $\beta$  type II receptor (TBR-II), human TGF- $\beta$  type I receptor (ALK-5), human activin receptor type IA (ALK-2), and type IB (ALK-4), ALKs 1 & 3 and mouse ALK-6.

Figure 4 shows, schematically, the structures for Daf-1, Act R-II, Act R-IIB, TBR-II, TBR-I/ALK-5, ALK's -1, -2 (Act RIA), -3, -4 (Act RIB) & -6.

Figure 5 shows the sequence alignment of the cysteine-rich domains of the ALKs, TBR-II, Act R-II, Act R-IIB and daf-1 receptors.

Figure 6 is a comparison of kinase domains of serine/threonine kinases, showing the percentage amino-acid identity of the kinase domains.

Figure 7 shows the pairwise alignment relationship between the kinase domains of the receptor serine/threonine kinases. The dendrogram was generated using the Jotun-Hein alignment program (Hein (1990) Meth. Enzymol. 183, 626-645).

#### Brief Description of the Sequence Listings

Sequences 1 and 2 are the nucleotide and deduced amino-acid sequences of cDNA for hALK-1 (clone HP57).

Sequences 3 and 4 are the nucleotide and deduced amino-acid sequences of cDNA for hALK-2 (clone HP53).

Sequences 5 and 6 are the nucleotide and deduced amino-acid sequences of cDNA for hALK-3 (clone ONF5).

Sequences 7 and 8 the nucleotide and deduced amino-acid sequences of cDNA for hALK-4 (clone 11H8), complemented with PCR product encoding extracellular domain.

Sequences 9 and 10 are the nucleotide and deduced amino-acid sequences of cDNA for hALK-5 (clone EMBLA).

Sequences 11 and 12 are the nucleotide and deduced amino-acid sequences of cDNA for mALK-1 (clone AM6).

Sequences 13 and 14 are the nucleotide and deduced amino-acid sequences of cDNA for mALK-3 (clones ME-7 and ME-D).

5 Sequences 15 and 16 are the nucleotide and deduced amino-acid sequences of cDNA for mALK-4 (clone 8a1).

Sequences 17 and 18 are the nucleotide and deduced amino-acid sequences of cDNA for mALK-6 (clone ME-6).

10 Sequence 19 (B1-S) is a sense primer, extracellular domain, cysteine-rich region, BamHI site at 5' end, 28-mer, 64-fold degeneracy.

Sequence 20 (B3-S) is a sense primer, kinase domain II, BamHI site at 5' end, 25-mer, 162-fold degeneracy.

15 Sequence 21 (B7-S) is a sense primer, kinase domain VIB, S/T kinase specific residues, BamHI site at 5' end, 24-mer, 288-fold degeneracy.

Sequence 22 (E8-AS) is an anti-sense primer, kinase domain, S/T kinase-specific residues EcoRI site at 5' end, 20-mer, 18-fold degeneracy.

20 ~~Sequence 23 is an oligonucleotide probe.~~

Sequence 24 is a 5' primer.

Sequence 25 is a 3' primer.

Sequence 26 is a consensus sequence in Subdomain I.

Sequences 27 and 28 are novel sequence motifs in Subdomain VIB.

25 Sequence 29 is a novel sequence motif in Subdomain VIII.

#### Description of the Invention

30 As described in more detail below, nucleic acid sequences have been isolated, coding for a new sub-family of serine/threonine receptor kinases. The term nucleic acid molecules as used herein refers to any sequence which codes for the murine, human or mammalian form, amino-acid sequences of which are presented herein. It is understood that the well known phenomenon of codon degeneracy provides  
35 for a great deal of sequence variation and all such varieties are included within the scope of this invention.



The nucleic acid sequences described herein may be used to clone the respective genomic DNA sequences in order to study the genes' structure and regulation. The murine and human cDNA or genomic sequences can also be used to  
5 isolate the homologous genes from other mammalian species. The mammalian DNA sequences can be used to study the receptors' functions in various in vitro and in vivo model systems.

As exemplified below for ALK-5 cDNA, it is also  
10 recognised that, given the sequence information provided herein, the artisan could easily combine the molecules with a pertinent promoter in a vector, so as to produce a cloning vehicle for expression of the molecule. The promoter and coding molecule must be operably linked via  
15 any of the well-recognized and easily-practised methodologies for so doing. The resulting vectors, as well as the isolated nucleic acid molecules themselves, may be used to transform prokaryotic cells (e.g. E. coli), or  
transfect eukaryotes such as yeast (S. cerevisiae), PAE,  
20 COS or CHO cell lines. Other appropriate expression systems will also be apparent to the skilled artisan.

Several methods may be used to isolate the ligands for the ALKs. As shown for ALK-5 cDNA, cDNA clones encoding the active open reading frames can be subcloned into  
25 expression vectors and transfected into eukaryotic cells, for example COS cells. The transfected cells which can express the receptor can be subjected to binding assays for radioactively-labelled members of the TGF- $\beta$  superfamily (TGF- $\beta$ , activins, inhibins, bone morphogenic proteins and  
30 müllerian-inhibiting substances), as it may be expected that the receptors will bind members of the TGF- $\beta$  superfamily. Various biochemical or cell-based assays can be designed to identify the ligands, in tissue extracts or conditioned media, for receptors in which a ligand is not  
35 known. Antibodies raised to the receptors may also be used to identify the ligands, using the immunoprecipitation of the cross-linked complexes. Alternatively, purified

receptor could be used to isolate the ligands using an affinity-based approach. The determination of the expression patterns of the receptors may also aid in the isolation of the ligand. These studies may be carried out using ALK DNA or RNA sequences as probes to perform in situ hybridisation studies.

The use of various model systems or structural studies should enable the rational development of specific agonists and antagonists useful in regulating receptor function. It may be envisaged that these can be peptides, mutated ligands, antibodies or other molecules able to interact with the receptors.

The foregoing provides examples of the invention Applicants intend to claim which includes, inter alia, isolated nucleic acid molecules coding for activin receptor-like kinases (ALKs), as defined herein. These include such sequences isolated from mammalian species such as mouse, human, rat, rabbit and monkey.

~~The following description relates to specific~~ embodiments. It will be understood that the specification and examples are illustrative but not limitative of the present invention and that other embodiments within the spirit and scope of the invention will suggest themselves to those skilled in the art.

#### Preparation of mRNA and Construction of a cDNA Library

For construction of a cDNA library, poly (A)<sup>+</sup> RNA was isolated from a human erythroleukemia cell line (HEL 92.1.7) obtained from the American Type Culture Collection (ATCC TIB 180). These cells were chosen as they have been shown to respond to both activin and TGF- $\beta$ . Moreover leukaemic cells have proved to be rich sources for the cloning of novel receptor tyrosine kinases (Partanen et al (1990) Proc. Natl. Acad. Sci. USA 87, 8913-8917 and (1992) Mol. Cell. Biol. 12, 1698-1707). (Total) RNA was prepared by the guanidinium isothiocyanate method (Chirgwin et al (1979) Biochemistry 18, 5294-5299). mRNA was selected using the poly-A or poly AT tract mRNA isolation kit

(Promega, Madison, Wisconsin, U.S.A.) as described by the manufacturers, or purified through an oligo (dT)-cellulose column as described by Aviv and Leder (1972) Proc. Natl. Acad. Sci. USA 69, 1408-1412. The isolated mRNA was used for the synthesis of random primed (Amersham) cDNA, that was used to make a  $\lambda$ gt10 library with  $1 \times 10^5$  independent cDNA clones using the Riboclone cDNA synthesis system (Promega) and  $\lambda$ gt10 in vitro packaging kit (Amersham) according to the manufacturers' procedures. An amplified oligo (dT) primed human placenta  $\lambda$ ZAPII cDNA library of  $5 \times 10^5$  independent clones was used. Poly (A)<sup>+</sup> RNA isolated from AG1518 human foreskin fibroblasts was used to prepare a primary random primed  $\lambda$ ZAPII cDNA library of  $1.5 \times 10^6$  independent clones using the RiboClone cDNA synthesis system and Gigapack Gold II packaging extract (Stratagene). In addition, a primary oligo (dT) primed human foreskin fibroblast  $\lambda$ gt10 cDNA library (Claesson-Welsh et al (1989) Proc. Natl. Acad. Sci. USA. 86 4917-4912) was prepared. An amplified oligo (dT) primed HEL cell  $\lambda$ gt11 cDNA library of  $1.5 \times 10^6$  independent clones (Poncz et al (1987) Blood 69 219-223) was used. A twelve-day mouse embryo  $\lambda$ EXIox cDNA library was obtained from Novagen (Madison, Wisconsin, U.S.A.); a mouse placenta  $\lambda$ ZAPII cDNA library was also used.

#### 25 Generation of cDNA Probes by PCR

For the generation of cDNA probes by PCR (Lee et al (1988) Science 239, 1288-1291) degenerate PCR primers were constructed based upon the amino-acid sequence similarity between the mouse activin type II receptor (Mathews and Vale (1991) Cell 65, 973-982) and daf-1 (George et al (1990) Cell 61, 635-645) in the kinase domains II and VIII. Figure 1 shows the aligned serine/threonine kinase domains (I-VIII), of four related receptors of the TGF- $\beta$  superfamily, i.e. hTBR-II, mActR-IIB, mActR-II and the daf-1 gene product, using the nomenclature of the subdomains according to Hanks et al (1988) Science 241, 45-52.

Several considerations were applied in the design of the PCR primers. The sequences were taken from regions of homology between the activin type II receptor and the daf-1 gene product, with particular emphasis on residues that confer serine/threonine specificity (see Table 2) and on residues that are shared by transmembrane kinase proteins and not by cytoplasmic kinases. The primers were designed so that each primer of a PCR set had an approximately similar GC composition, and so that self complementarity and complementarity between the 3' ends of the primer sets were avoided. Degeneracy of the primers was kept as low as possible, in particular avoiding serine, leucine and arginine residues (6 possible codons), and human codon preference was applied. Degeneracy was particularly avoided at the 3' end as, unlike the 5' end, where mismatches are tolerated, mismatches at the 3' end dramatically reduce the efficiency of PCR.

In order to facilitate directional subcloning, restriction enzyme sites were included at the 5' end of the primers, with a GC clamp, which permits efficient restriction enzyme digestion. The primers utilised are shown in Figure 2. Oligonucleotides were synthesized using Gene assembler plus (Pharmacia - LKB) according to the manufacturers instructions.

The mRNA prepared from HEL cells as described above was reverse-transcribed into cDNA in the presence of 50 mM Tris-HCl, pH 8.3, 8 mM MgCl<sub>2</sub>, 30 mM KCl, 10 mM dithiothreitol, 2mM nucleotide triphosphates, excess oligo (dT) primers and 34 units of AMV reverse transcriptase at 42°C for 2 hours in 40 µl of reaction volume. Amplification by PCR was carried out with a 7.5% aliquot (3 µl) of the reverse-transcribed mRNA, in the presence of 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 M MgCl<sub>2</sub>, 0.01% gelatin, 0.2 mM nucleotide triphosphates, 1 µM of both sense and antisense primers and 2.5 units of Taq polymerase (Perkin Elmer Cetus) in 100 µl reaction volume. Amplifications were performed on a thermal cycler (Perkin Elmer Cetus)

using the following program: first 5 thermal cycles with denaturation for 1 minute at 94°C, annealing for 1 minute at 50°C, a 2 minute ramp to 55°C and elongation for 1 minute at 72°C, followed by 20 cycles of 1 minute at 94°C, 30 seconds at 55°C and 1 minute at 72°C. A second round of PCR was performed with 3 µl of the first reaction as a template. This involved 25 thermal cycles, each composed of 94°C (1 min), 55°C (0.5 min), 72°C (1 min).

General procedures such as purification of nucleic acids, restriction enzyme digestion, gel electrophoresis, transfer of nucleic acid to solid supports and subcloning were performed essentially according to established procedures as described by Sambrook *et al*, (1989), Molecular cloning: A Laboratory Manual, 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory (Cold Spring Harbor, New York, USA).

Samples of the PCR products were digested with BamHI and EcoRI and subsequently fractionated by low melting point agarose gel electrophoresis. Bands corresponding to the approximate expected sizes, (see Table 1: ≈460 bp for primer pair B3-S and E8-AS and ≈ 140 bp for primer pair B7-S and E8-AS) were excised from the gel and the DNA was purified. Subsequently, these fragments were ligated into pUC19 (Yanisch-Perron *et al* (1985) Gene 33, 103-119), which had been previously linearised with BamHI and EcoRI and transformed into E. coli strain DH5α using standard protocols (Sambrook *et al*, *supra*). Individual clones were sequenced using standard double-stranded sequencing techniques and the dideoxynucleotide chain termination method as described by Sanger *et al* (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467, and T7 DNA polymerase.

Employing Reverse Transcriptase PCR on HEL mRNA with the primer pair B3-S and E8-AS, three PCR products were obtained, termed 11.1, 11.2 and 11.3, that corresponded to novel genes. Using the primer pair B7-S and E8-AS, an additional novel PCR product was obtained termed 5.2.

TABLE 1

NAME OF PCR PRODUCT	PRIMERS	INSERT SIZE (bp)	SIZE OF DNA FRAGMENT IN mActRII/hTBRII CLONES (bp)	SEQUENCE IDENTITY WITH SEQUENCE mActRII/hTBRII (%)	SEQUENCE IDENTITY BETWEEN mActRII and TBR-II (%)
11.1	B3-S/E8-AS	460	460	46/40	42
11.2	B3-S/E8-AS	460	460	49/44	47
11.3	B3-S/E8-AS	460	460	44/36	48
11.29	B3-S/E8-AS	460	460	ND/100	ND
9.2	B1-S/E8-AS	800	795	100/ND	ND
5.2	B7-S/E8-AS	140	143	40/38	60

#### 15 Isolation of cDNA Clones

The PCR products obtained were used to screen various cDNA libraries described supra. Labelling of the inserts of PCR products was performed using random priming method (Feinberg and Vogelstein (1983) Anal. Biochem, 132 6-13) using the Megaprime DNA labelling system (Amersham). The oligonucleotide derived from the sequence of the PCR product 5.2 was labelled by phosphorylation with T4 polynucleotide kinase following standard protocols (Sambrook et al, supra). Hybridization and purification of positive bacteriophages were performed using standard molecular biological techniques.

The double-stranded DNA clones were all sequenced using the dideoxynucleotide chain-termination method as described by Sanger et al, supra, using T7 DNA polymerase (Pharmacia - LKB) or Sequenase (U.S. Biochemical Corporation, Cleveland, Ohio, U.S.A.). Compressions of nucleotides were resolved using 7-deaza-GTP (U.S. Biochemical Corp.) DNA sequences were analyzed using the DNA STAR computer program (DNA STAR Ltd. U.K.). Analyses of the sequences obtained revealed the existence of six

distinct putative receptor serine/threonine kinases which have been named ALK 1-6.

To clone cDNA for ALK-1 the oligo (dT) primed human placenta cDNA library was screened with a radiolabelled insert derived from the PCR product 11.3; based upon their restriction enzyme digestion patterns, three different types of clones with approximate insert sizes of 1.7 kb, 2 kb & 3.5 kb were identified. The 2 kb clone, named HP57, was chosen as representative of this class and subjected to complete sequencing. Sequence analysis of ALK-1 revealed a sequence of 1984 nucleotides including a poly-A tail (SEQ ID No. 1). The longest open reading frame encodes a protein of 503 amino-acids, with high sequence similarity to receptor serine/threonine kinases (see below). The first methionine codon, the putative translation start site, is at nucleotide 283-285 and is preceded by an in-frame stop codon. This first ATG is in a more favourable context for translation initiation (Kozak (1987) Nucl. Acids Res., 15, 8125-8148) than the second and third in-frame ATG at nucleotides 316-318 and 325-327. The putative initiation codon is preceded by a 5' untranslated sequence of 282 nucleotides that is GC-rich (80% GC), which is not uncommon for growth factor receptors (Kozak (1991) J. Cell Biol., 115, 887-903). The 3' untranslated sequence comprises 193 nucleotides and ends with a poly-A tail. No bona fide poly-A addition signal is found, but there is a sequence (AATACA), 17-22 nucleotides upstream of the poly-A tail, which may serve as a poly-A addition signal.

ALK-2 cDNA was cloned by screening an amplified oligo (dT) primed human placenta cDNA library with a radiolabelled insert derived from the PCR product 11.2. Two clones, termed HP53 and HP64, with insert sizes of 2.7 kb and 2.4 kb respectively, were identified and their sequences were determined. No sequence difference in the overlapping clones was found, suggesting they are both derived from transcripts of the same gene.

Sequence analysis of cDNA clone HP53 (SEQ ID No. 3) revealed a sequence of 2719 nucleotides with a poly-A tail. The longest open reading frame encodes a protein of 509 amino-acids. The first ATG at nucleotides 104-106 agrees favourably with Kozak's consensus sequence with an A at position 3. This ATG is preceded in-frame by a stop codon. There are four ATG codons in close proximity further downstream, which agree with the Kozak's consensus sequence (Kozak, supra), but according to Kozak's scanning model the first ATG is predicted to be the translation start site. The 5' untranslated sequence is 103 nucleotides. The 3' untranslated sequence of 1089 nucleotides contains a polyadenylation signal located 9-14 nucleotides upstream from the poly-A tail. The cDNA clone HP64 lacks 498 nucleotides from the 5' end compared to HP53, but the sequence extended at the 3' end with 190 nucleotides and poly-A tail is absent. This suggests that different polyadenylation sites occur for ALK-2. In Northern blots, however, only one transcript was detected (see below).

The cDNA for human ALK-3 was cloned by initially screening an oligo (dT) primed human foreskin fibroblast cDNA library with an oligonucleotide (SEQ ID No. 23) derived from the PCR product 5.2. One positive cDNA clone with an insert size of 3 kb, termed ON11, was identified. However, upon partial sequencing, it appeared that this clone was incomplete; it encodes only part of the kinase domain and lacks the extracellular domain. The most 5' sequence of ON11, a 540 nucleotide XbaI restriction fragment encoding a truncated kinase domain, was subsequently used to probe a random primed fibroblast cDNA library from which one cDNA clone with an insert size of 3 kb, termed ONF5, was isolated (SEQ ID No. 5). Sequence analysis of ONF5 revealed a sequence of 2932 nucleotides without a poly-A tail, suggesting that this clone was derived by internal priming. The longest open reading frame codes for a protein of 532 amino-acids. The first ATG codon which is compatible with Kozak's consensus



sequence (Kozak, supra), is at 310-312 nucleotides and is preceded by an in-frame stop codon. The 5' and 3' untranslated sequences are 309 and 1027 nucleotides long, respectively.

5        ALK-4 cDNA was identified by screening a human oligo (dT) primed human erythroleukemia cDNA library with the radiolabelled insert of the PCR product 11.1 as a probe. One cDNA clone, termed 11H8, was identified with an insert size of 2 kb (SEQ ID No. 7). An open reading frame was  
10 found encoding a protein sequence of 383 amino-acids encoding a truncated extracellular domain with high similarity to receptor serine/threonine kinases. The 3' untranslated sequence is 818 nucleotides and does not contain a poly-A tail, suggesting that the cDNA was  
15 internally primed. cDNA encoding the complete extracellular domain (nucleotides 1-366) was obtained from HEL cells by RT-PCR with 5' primer (SEQ ID No. 24) derived in part from sequence at translation start site of SKR-2 (a cDNA sequence deposited in GenBank data base, accession  
20 number L10125, that is identical in part to ALK-4) and 3' primer (SEQ ID No. 25) derived from 11H8 cDNA clone.

      ALK-5 was identified by screening the random primed HEL cell  $\lambda$ gt 10 cDNA library with the PCR product 11.1 as a probe. This yielded one positive clone termed EMBLA  
25 (insert size of 5.3 kb with 2 internal EcoRI sites). Nucleotide sequencing revealed an open reading frame of 1509 bp, coding for 503 amino-acids. The open reading frame was flanked by a 5' untranslated sequence of 76 bp, and a 3' untranslated sequence of 3.7 kb which was not  
30 completely sequenced. The nucleotide and deduced amino-acid sequences of ALK-5 are shown in SEQ ID Nos. 9 and 10. In the 5' part of the open reading frame, only one ATG codon was found; this codon fulfils the rules of translation initiation (Kozak, supra). An in-frame stop  
35 codon was found at nucleotides (-54)-(-52) in the 5' untranslated region. The predicted ATG start codon is followed by a stretch of hydrophobic amino-acid residues

which has characteristics of a cleavable signal sequence. Therefore, the first ATG codon is likely to be used as a translation initiation site. A preferred cleavage site for the signal peptidase, according to von Heijne (1986) Nucl. Acid. Res. 14, 4683-4690, is located between amino-acid residues 24 and 25. The calculated molecular mass of the primary translated product of the ALK-5 without signal sequence is 53,646 Da.

Screening of the mouse embryo  $\lambda$ EX Iox cDNA library using PCR, product 11.1 as a probe yielded 20 positive clones. DNAs from the positive clones obtained from this library were digested with EcoRI and HindIII, electrophoretically separated on a 1.3% agarose gel and transferred to nitrocellulose filters according to established procedures as described by Sambrook et al, supra. The filters were then hybridized with specific probes for human ALK-1 (nucleotide 288-670), ALK-2 (nucleotide 1-581), ALK-3 (nucleotide 79-824) or ALK-4 (nucleotide 1178-1967). Such analyses revealed that a clone termed ME-7 hybridised with the human ALK-3 probe. However, nucleotide sequencing revealed that this clone was incomplete, and lacked the 5' part of the translated region. Screening the same cDNA library with a probe corresponding to the extracellular domain of human ALK-3 (nucleotides 79-824) revealed the clone ME-D. This clone was isolated and the sequence was analyzed. Although this clone was incomplete in the 3' end of the translated region, ME-7 and ME-D overlapped and together covered the complete sequence of mouse ALK-3. The predicted amino-acid sequence of mouse ALK-3 is very similar to the human sequence; only 8 amino-acid residues differ (98% identity; see SEQ ID No. 14) and the calculated molecular mass of the primary translated product without the putative signal sequence is 57,447 Da.

Of the clones obtained from the initial library screening with PCR product 11.1, four clones hybridized to the probe corresponding to the conserved kinase domain of

ALK-4 but not to probes from more divergent parts of ALK-1 to -4. Analysis of these clones revealed that they have an identical sequence which differs from those of ALK-1 to -5 and was termed ALK-6. The longest clone ME6 with a 2.0 kb insert was completely sequenced yielding a 1952 bp fragment consisting of an open reading frame of 1506 bp (502 amino-acids), flanked by a 5' untranslated sequence of 186 bp, and a 3' untranslated sequence of 160 bp. The nucleotide and predicted amino-acid sequences of mouse ALK-6 are shown in SEQ ID Nos. 17 and 18. No polyadenylation signal was found in the 3' untranslated region of ME6, indicating that the cDNA was internally primed in the 3' end. Only one ATG codon was found in the 5' part of the open reading frame, which fulfils the rules for translation initiation (Kozak, supra), and was preceded by an in-frame stop codon at nucleotides 163-165. However, a typical hydrophobic leader sequence was not observed at the N terminus of the translated region. Since there is no ATG codon and putative hydrophobic leader sequence, this ATG codon is likely to be used as a translation initiation site. The calculated molecular mass of the primary translated product with the putative signal sequence is 55,576 Da.

Mouse ALK-1 (clone AM6 with 1.9 kb insert) was obtained from the mouse placenta  $\lambda$ ZAPII cDNA library using human ALK-1 cDNA as a probe (see SEQ ID No. 11). Mouse ALK-4 (clone 8a1 with 2.3kb insert) was also obtained from this library using human ALK-4 cDNA library as a probe (SEQ ID No. 15).

To summarise, clones HP22, HP57, ONF1, ONF3, ONF4 and HP29 encode the same gene, ALK-1. Clone AM6 encodes mouse ALK-1. HP53, HP64 and HP84 encode the same gene, ALK-2. ONF5, ONF2 and ON11 encode the same gene ALK-3. ME-7 and ME-D encode the mouse counterpart of human ALK-3. 11H8 encodes a different gene ALK-4, whilst 8a1 encodes the mouse equivalent. EMBLA encodes ALK-5, and ME-6 encodes ALK-6.

The sequence alignment between the 6 ALK genes and TBR-II, mActR-II and ActR-IIB is shown in Figure 3. These molecules have a similar domain structure; an N-terminal predicted hydrophobic signal sequence (von Heijne (1986) Nucl. Acids Res. 14: 4683-4690) is followed by a relatively small extracellular cysteine-rich ligand binding domain, a single hydrophobic transmembrane region (Kyte & Doolittle (1982) J. Mol. Biol. 157, 105-132) and a C-terminal intracellular portion, which consists almost entirely of a kinase domain (Figures 3 and 4).

The extracellular domains of these receptors have cysteine-rich regions, but they show little sequence similarity; for example, less than 20% sequence identity is found between Daf-1, ActR-II, TBR-II and ALK-5. The ALKs appear to form a subfamily as they show higher sequence similarities (15-47% identity) in their extracellular domains. The extracellular domains of ALK-5 and ALK-4 have about 29% sequence identity. In addition, ALK-3 and ALK-6 share a high degree of sequence similarity in their extracellular domains (46% identity).

The positions of many of the cysteine residues in all receptors can be aligned, suggesting that the extracellular domains may adopt a similar structural configuration. See Figure 5 for ALKs-1,-2,-3 & 5. Each of the ALKs (except ALK-6) has a potential N-linked glycosylation site, the position of which is conserved between ALK-1 and ALK-2, and between ALK-3, ALK-4 and ALK-5 (see Figure 4).

The sequence similarities in the kinase domains between daf-1, ActR-II, TBR-II and ALK-5 are approximately 40%, whereas the sequence similarity between the ALKs 1 to 6 is higher (between 59% and 90%; see Figure 6). Pairwise comparison using the Jutun-Hein sequence alignment program (Hein (1990) Meth, Enzymol., 183, 626-645), between all family members, identifies the ALKs as a separate subclass among serine/threonine kinases (Figure 7).

The catalytic domains of kinases can be divided into 12 subdomains with stretches of conserved amino-acid

residues. The key motifs are found in serine/threonine kinase receptors suggesting that they are functional kinases. The consensus sequence for the binding of ATP (Gly-X-Gly-X-X-Gly in subdomain I followed by a Lys residue further downstream in subdomain II) is found in all the ALKs.

The kinase domains of daf-1, ActR-II, and ALKs show approximately equal sequence similarity with tyrosine and serine/threonine protein kinases. However analysis of the amino-acid sequences in subdomains VI and VIII, which are the most useful to distinguish a specificity for phosphorylation of tyrosine residues versus serine/threonine residues (Hanks et al (1988) Science 241 42-52) indicates that these kinases are serine/threonine kinases; refer to Table 2.

TABLE 2

KINASE	SUBDOMAINS	
	VIB	VIII
Serine/threonine kinase consensus	DLKPEN	G (T/S) XX (Y/F) X
5 Tyrosine kinase consensus	DLAARN	XP(I/V) (K/R) W (T/M)
Act R-II	DIKSKN	GTRRYM
Act R-IIB	DFKSKN	GTRRYM
TBR-II	DLKSSN	GTARYM
ALK-I	DFKSRN	GTKRYM
10 ALK -2, -3, -4, -5, & -6	DLKSKN	GTKRYM

The sequence motifs DLKSKN (Subdomain VIB) and GTKRYM (Subdomain VIII), that are found in most of the serine/threonine kinase receptors, agree well with the consensus sequences for all protein serine/threonine kinase receptors in these regions. In addition, these receptors, except for ALK-1, do not have a tyrosine residue surrounded by acidic residues between subdomains VII and VIII, which is common for tyrosine kinases. A unique characteristic of the members of the ALK serine/threonine kinase receptor family is the presence of two short inserts in the kinase

domain between subdomains VIA and VIB and between subdomains X and XI. In the intracellular domain, these regions, together with the juxtamembrane part and C-terminal tail, are the most divergent between family members (see Figures 3 and 4). Based on the sequence similarity with the type II receptors for TGF- $\beta$  and activin, the C termini of the kinase domains of ALKs -1 to -6 are set at Ser-495, Ser-501, Ser-527, Gln-500, Gln-498 and Ser-497, respectively.

#### 10 mRNA Expression

The distribution of ALK-1, -2, -3, -4 was determined by Northern blot analysis. A Northern blot filter with mRNAs from different human tissues was obtained from Clontech (Palo Alto, C.A.). The filters were hybridized with  $^{32}\text{P}$ -labelled probes at 42°C overnight in 50% formaldehyde, 5 x standard saline citrate (SSC; 1xSSC is 50mM sodium citrate, pH 7.0, 150 mM NaCl), 0.1% SDS, 50 mM sodium phosphate, 5 x Denhardt's solution and 0.1 mg/ml salmon sperm DNA. In order to minimize cross-hybridization, probes were used that did not encode part of the kinase domains, but corresponded to the highly diverged sequences of either 5' untranslated and ligand-binding regions (probes for ALK-1, -2 and -3) or 3' untranslated sequences (probe for ALK-4). The probes were labelled by random priming using the Multiprime (or Mega-prime) DNA labelling system and [ $\alpha$ - $^{32}\text{P}$ ] dCTP (Feinberg & Vogelstein (1983) Anal. Biochem. 132: 6-13). Unincorporated label was removed by Sephadex G-25 chromatography. Filters were washed at 65°C, twice for 30 minutes in 2.5 x SSC, 0.1% SDS and twice for 30 minutes in 0.3 x SSC, 0.1% SDS before being exposed to X-ray film. Stripping of blots was performed by incubation at 90-100°C in water for 20 minutes.

The ALK-5 mRNA size and distribution were determined by Northern blot analysis as above. An EcoRI fragment of 980bp of the full length ALK-5 cDNA clone, corresponding to the C-terminal part of the kinase domain and 3'

untranslated region (nucleotides 1259-2232 in SEQ ID No. 9) was used as a probe. The filter was washed twice in 0.5 x SSC, 0.1% SDS at 55°C for 15 minutes.

Using the probe for ALK-1, two transcripts of 2.2 and 4.9kb were detected. The ALK-1 expression level varied strongly between different tissues, high in placenta and lung, moderate in heart, muscle and kidney, and low (to not detectable) in brain, liver and pancreas. The relative ratios between the two transcripts were similar in most tissues; in kidney, however, there was relatively more of the 4.9 kb transcript. By reprobing the blot with a probe for ALK-2, one transcript of 4.0 kb was detected with a ubiquitous expression pattern. Expression was detected in every tissue investigated and was highest in placenta and skeletal muscle. Subsequently the blot was reprobed for ALK-3. One major transcript of 4.4 kb and a minor transcript of 7.9 kb were detected. Expression was high in skeletal muscle, in which also an additional minor transcript of 10 kb was observed. Moderate levels of ALK-3 mRNA were detected in heart, placenta, kidney and pancreas, and low (to not detectable) expression was found in brain, lung and liver. The relative ratios between the different transcripts were similar in the tested tissues, the 4.4 kb transcript being the predominant one, with the exception for brain where both transcripts were expressed at a similar level. Probing the blot with ALK-4 indicated the presence of a transcript with the estimated size of 5.2 kb and revealed an ubiquitous expression pattern. The results of Northern blot analysis using the probe for ALK-5 showed that a 5.5 kb transcript is expressed in all human tissues tested, being most abundant in placenta and least abundant in brain and heart.

The distribution of mRNA for mouse ALK-3 and -6 in various mouse tissues was also determined by Northern blot analysis. A multiple mouse tissue blot was obtained from Clontech, Palo Alto, California, U.S.A. The filter was hybridized as described above with probes for mouse ALK-3



and ALK-6. The EcoRI-PstI restriction fragment, corresponding to nucleotides 79-1100 of ALK-3, and the SacI-HpaI fragment, corresponding to nucleotides 57-720 of ALK-6, were used as probes. The filter was washed at 65°C  
5 twice for 30 minutes in 2.5 x SSC, 0.1% SDS and twice for 30 minutes with 0.3 x SSC, 0.1% SDS and then subjected to autoradiography.

Using the probe for mouse ALK-3, a 1.1 kb transcript was found only in spleen. By reprobing the blot with the  
10 ALK-6 specific probe, a transcript of 7.2 kb was found in brain and a weak signal was also seen in lung. No other signal was seen in the other tissues tested, i.e. heart, liver, skeletal muscle, kidney and testis.

All detected transcript sizes were different, and thus  
15 no cross-reaction between mRNAs for the different ALKs was observed when the specific probes were used. This suggests that the multiple transcripts of ALK-1 and ALK-3 are coded from the same gene. The mechanism for generation of the different transcripts is unknown at present; they may be  
20 formed by alternative mRNA splicing, differential polyadenylation, use of different promoters, or by a combination of these events. Differences in mRNA splicing in the regions coding for the extracellular domains may lead to the synthesis of receptors with different  
25 affinities for ligands, as was shown for mActR-IIB (Attisano et al (1992) Cell 68, 97-108) or to the production of soluble binding protein.

~~-----~~The above experiments describe the isolation of nucleic acid sequences coding for new family of human  
30 receptor kinases. The cDNA for ALK-5 was then used to determine the encoded protein size and binding properties.  
Properties of the ALKs cDNA Encoded Proteins

To study the properties of the proteins encoded by the different ALK cDNAs, the cDNA for each ALK was subcloned  
35 into a eukaryotic expression vector and transfected into various cell types and then subjected to immunoprecipitation using a rabbit antiserum raised against

a synthetic peptide corresponding to part of the intracellular juxtamembrane region. This region is divergent in sequence between the various serine/threonine kinase receptors. The following amino-acid residues were used:

5	ALK-1	145-166
	ALK-2	151-172
	ALK-3	181-202
	ALK-4	153-171
10	ALK-5	158-179
	ALK-6	151-168

The rabbit antiserum against ALK-5 was designated VPN.

The peptides were synthesized with an Applied Biosystems 430A Peptide Synthesizer using t-butoxycarbonyl chemistry and purified by reversed-phase high performance liquid chromatography. The peptides were coupled to keyhole limpet haemocyanin (Calbiochem-Behring) using glutaraldehyde, as described by Guilleck *et al* (1985) EMBO J. 4, 2869-2877. The coupled peptides were mixed with Freund's adjuvant and used to immunize rabbits.

#### Transient transfection of the ALK-5 cDNA

COS-1 cells (American Type Culture Collection) and the R mutant of Mv1Lu cells (for references, see below) were cultured in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum (FBS) and 100 units/ml penicillin and 50 µg/ml streptomycin in 5% CO<sub>2</sub> atmosphere at 37°C. The ALK-5 cDNA (nucleotides (-76) - 2232), which includes the complete coding region, was cloned in the pSV7d vector (Truett *et al*, (1985) DNA 4, 333-349), and used for transfection. Transfection into COS-1 cells was performed by the calcium phosphate precipitation method (Wigler *et al* (1979) Cell 16, 777-785). Briefly, cells were seeded into 6-well cell culture plates at a density of 5x10<sup>5</sup> cells/well, and transfected the following day with 10 µg of recombinant plasmid. After overnight incubation, cells were washed three times with a buffer containing 25 mM Tris-HCl, pH 7.4, 138 mM NaCl, 5 mM KCl, 0.7 mM CaCl<sub>2</sub>, 0.5

mM  $\text{MgCl}_2$  and 0.6 mM  $\text{Na}_2\text{HPO}_4$ , and then incubated with Dulbecco's modified Eagle's medium containing FBS and antibiotics. Two days after transfection, the cells were metabolically labelled by incubating the cells for 6 hours  
5 in methionine and cysteine-free MCDB 104 medium with 150  $\mu\text{Ci/ml}$  of [ $^{35}\text{S}$ ]-methionine and [ $^{35}\text{S}$ ]-cysteine (in vivo labelling mix; Amersham). After labelling, the cells were washed with 150 mM NaCl, 25 mM Tris-HCl, pH 7.4, and then solubilized with a buffer containing 20mM Tris-HCl, pH 7.4,  
10 150 mM NaCl, 10 mM EDTA, 1% Triton X-100, 1% deoxycholate, 1.5% Trasylol (Bayer) and 1 mM phenylmethylsulfonylfluoride (PMSF; Sigma). After 15 minutes on ice, the cell lysates were pelleted by centrifugation, and the supernatants were then incubated with 7  $\mu\text{l}$  of preimmune serum for 1.5 hours  
15 at 4°C. Samples were then given 50  $\mu\text{l}$  of protein A-Sepharose (Pharmacia-LKB) slurry (50% packed beads in 150 mM NaCl, 20 mM Tris-HCl, pH 7.4, 0.2% Triton X100) and incubated for 45 minutes at 4°C. The beads were spun down by centrifugation, and the supernatants (1 ml) were then  
20 incubated with either 7  $\mu\text{l}$  of preimmune serum or the VPN antiserum for 1.5 hours at 4°C. For blocking, 10  $\mu\text{g}$  of peptide was added together with the antiserum. Immune complexes were then given 50  $\mu\text{l}$  of protein A-Sepharose (Pharmacia - LKB) slurry (50% packed beads in 150 mM NaCl,  
25 20mM Tris-HCl, pH 7.4, 0.2% Triton X-100) and incubated for 45 minutes at 4°C. The beads were spun down and washed four times with a washing buffer (20 mM Tris-HCl, pH 7.4, 500 mM NaCl, 1% Triton X-100, 1% deoxycholate and 0.2% SDS), followed by one wash in distilled water. The immune  
30 complexes were eluted by boiling for 5 minutes in the SDS-sample buffer (100 mM Tris-HCl, pH 8.8, 0.01% bromophenol blue, 36% glycerol, 4% SDS) in the presence of 10 mM DTT, and analyzed by SDS-gel electrophoresis using 7-15% polyacrylamide gels (Blobel and Dobberstein, (1975) J. Cell  
35 Biol. 67, 835-851). Gels were fixed, incubated with Amplify (Amersham) for 20 minutes, and subjected to fluorography. A component of 53Da was seen. This

component was not seen when preimmune serum was used, or when 10 µg blocking peptide was added together with the antiserum. Moreover, it was not detectable in samples derived from untransfected COS-1 cells using either  
5 preimmune serum or the antiserum.

#### Digestion with Endoglycosidase F

Samples immunoprecipitated with the VPN antisera obtained as described above were incubated with 0.5 U of endoglycosidase F (Boehringer Mannheim Biochemica) in a  
10 buffer containing 100 mM sodium phosphate, pH 6.1, 50 mM EDTA, 1% Triton X-100, 0.1% SDS and 1% β-mercaptoethanol at 37°C for 24 hours. Samples were eluted by boiling for 5 minutes in the SDS-sample buffer, and analyzed by SDS-polyacrylamide gel electrophoresis as described above.  
15 Hydrolysis of N-linked carbohydrates by endoglycosidase F shifted the 53 kDa band to 51 kDa. The extracellular domain of ALK-5 contains one potential acceptor site for N-glycosylation and the size of the deglycosylated protein is close to the predicted size of the core protein.

#### 20 Establishment of PAE Cell Lines Expressing ALK-5

In order to investigate whether the ALK-5 cDNA encodes a receptor for TGF-β, porcine aortic endothelial (PAE) cells were transfected with an expression vector containing the ALK-5 cDNA, and analyzed for the binding of <sup>125</sup>I-TGF-β1.  
25 PAE cells were cultured in Ham's F-12 medium supplemented with 10% FBS and antibiotics (Miyazono *et al.*, (1988) J. Biol. Chem. 263, 6407-6415). The ALK-5 cDNA was cloned into the cytomegalovirus (CMV)-based expression vector pcDNA I/NEO (Invitrogen), and transfected into PAE  
30 cells by electroporation. After 48 hours, selection was initiated by adding Geneticin (G418 sulphate; Gibco - BRL) to the culture medium at a final concentration of 0.5 mg/ml (Westermarck *et al.*, (1990) Proc. Natl. Acad. Sci. USA 87, 128-132). Several clones were obtained, and after analysis  
35 by immunoprecipitation using the VPN antiserum, one clone denoted PAE/TBR-1 was chosen and further analyzed.

### Iodination of TGF- $\beta$ 1, Binding and Affinity Crosslinking

Recombinant human TGF- $\beta$ 1 was iodinated using the chloramine T method according to Frolik *et al.*, (1984) J. Biol. Chem. 259, 10995-11000. Cross-linking experiments were performed as previously described (Ichijo *et al.*, (1990) Exp. Cell Res. 187, 263-269). Briefly, cells in 6-well plates were washed with binding buffer (phosphate-buffered saline containing 0.9 mM CaCl<sub>2</sub>, 0.49 mM MgCl<sub>2</sub> and 1 mg/ml bovine serum albumin (BSA)), and incubated on ice in the same buffer with <sup>125</sup>I-TGF- $\beta$ 1 in the presence or absence of excess unlabelled TGF- $\beta$ 1 for 3 hours. Cells were washed and cross-linking was done in the binding buffer without BSA together with 0.28 mM disuccinimidyl suberate (DSS; Pierce Chemical Co.) for 15 minutes on ice. The cells were harvested by the addition of 1 ml of detachment buffer (10 mM Tris-HCl, pH 7.4, 1 mM EDTA, 10% glycerol, 0.3 mM PMSF). The cells were pelleted by centrifugation, then resuspended in 50  $\mu$ l of solubilization buffer (125 mM NaCl, 10 mM Tris-HCl, pH 7.4, 1 mM EDTA, 1% Triton X-100, 0.3 mM PMSF, 1% Trasylol) and incubated for 40 minutes on ice. Cells were centrifuged again and supernatants were subjected to analysis by SDS-gel electrophoresis using 4-15% polyacrylamide gels, followed by autoradiography. <sup>125</sup>I-TGF- $\beta$ 1 formed a 70 kDa cross-linked complex in the transfected PAE cells (PAE/T $\beta$ R-I cells). The size of this complex was very similar to that of the TGF- $\beta$  type I receptor complex observed at lower amounts in the untransfected cells. A concomitant increase of 94 kDa TGF- $\beta$  type II receptor complex could also be observed in the PAE/T $\beta$ R-I cells. Components of 150-190 kDa, which may represent crosslinked complexes between the type I and type II receptors, were also observed in the PAE/T $\beta$ R-I cells.

In order to determine whether the cross-linked 70 kDa complex contained the protein encoded by the ALK-5 cDNA, the affinity cross-linking was followed by immunoprecipitation using the VPN antiserum. For this,

~~cells in 25 cm<sup>2</sup> flasks were used.~~ The supernatants obtained after cross-linking were incubated with 7  $\mu$ l of preimmune serum or VPN antiserum in the presence or absence of 10  $\mu$ g of peptide for 1.5h at 4°C. Immune complexes were then added to 50  $\mu$ l of protein A-Sepharose slurry and ~~incubated for 45 minutes at 4°C.~~ The protein A-Sepharose beads were washed four times with the washing buffer, once with distilled water, and the samples were analyzed by SDS-gel electrophoresis using 4-15% polyacrylamide gradient gels and autoradiography. A 70 kDa cross-linked complex was precipitated by the VPN antiserum in PAE/TBR-1 cells, and a weaker band of the same size was also seen in the untransfected cells, indicating that the untransfected PAE cells contained a low amount of endogenous ALK-5. The 70 kDa complex was not observed when preimmune serum was used, or when immune serum was blocked by 10  $\mu$ g of peptide. Moreover, a coprecipitated 94 kDa component could also be observed in the PAE/TBR-I cells. The latter component is ~~likely to represent a TGF- $\beta$  type II receptor complex,~~ since an antiserum, termed DRL, which was raised against a synthetic peptide from the C-terminal part of the TGF- $\beta$  type II receptor, precipitated a 94 kDa TGF- $\beta$  type II receptor complex, as well as a 70 kDa type I receptor complex from PAE/TBR-I cells.

The carbohydrate contents of ALK-5 and the TGF- $\beta$  type II receptor were characterized by deglycosylation using endoglycosidase F as described above and analyzed by SDS-polyacrylamide gel electrophoresis and autoradiography. The ALK-5 cross-linked complex shifted from 70 kDa to 66 kDa, whereas that of the type II receptor shifted from 94 kDa to 82 kDa. The observed larger shift of the type II receptor band compared with that of the ALK-5 band is consistent with the deglycosylation data of the type I and type II receptors on rat liver cells reported previously (Cheifetz et al (1988) J. Biol. Chem. 263, 16984-16991), and fits well with the fact that the porcine TGF- $\beta$  type II receptor has two N-glycosylation sites (Lin et al (1992)

Cell 68, 775-785), whereas ALK-5 has only one (see SEQ ID No. 9).

Binding of TGF- $\beta$ 1 to the type I receptor is known to be abolished by transient treatment of the cells with dithiothreitol (DTT) (Cheifetz and Massague (1991) J. Biol. Chem. 266, 20767-20772; Wrana *et al* (1992) Cell 71, 1003-1014). When analyzed by affinity cross-linking, binding of  $^{125}$ I-TGF- $\beta$ 1 to ALK-5, but not to the type II receptor, was completely abolished by DTT treatment of PAE/TBR-1 cells. Affinity cross-linking followed by immunoprecipitation by the VPN antiserum showed that neither the ALK-5 nor the type II receptor complexes was precipitated after DTT treatment, indicating that the VPN antiserum reacts only with ALK-5. The data show that the VPN antiserum recognizes a TGF- $\beta$  type I receptor, and that the type I and type II receptors form a heteromeric complex.

$^{125}$ I-TGF- $\beta$ 1 Binding & Affinity Crosslinking of Transfected COS Cells

Transient expression plasmids of ALKs -1 to -6 and TBR-II were generated by subcloning into the pSV7d expression vector or into the pcDNA I expression vector (Invitrogen). Transient transfection of COS-1 cells and iodination of TGF- $\beta$ 1 were carried out as described above. Crosslinking and immunoprecipitation were performed as described for PAE cells above.

Transfection of cDNAs for ALKs into COS-1 cells did not show any appreciable binding of  $^{125}$ I-TGF $\beta$ 1, consistent with the observation that type I receptors do not bind TGF- $\beta$  in the absence of type II receptors. When the TBR-II cDNA was co-transfected with cDNAs for the different ALKs, type I receptor-like complexes were seen, at different levels, in each case. COS-1 cells transfected with TBR-II and ALK cDNAs were analyzed by affinity crosslinking followed by immunoprecipitation using the DRL antisera or specific antisera against ALKs. Each one of the ALKs bound  $^{125}$ I-TGF- $\beta$ 1 and was coimmunoprecipitated with the TBR-II complex using the DRL antiserum. Comparison of the

efficiency of the different ALKs to form heteromeric complexes with TBR-II, revealed that ALK-5 formed such complexes more efficiently than the other ALKs. The size of the crosslinked complex was larger for ALK-3 than for other ALKs, consistent with its slightly larger size.

#### Expression of the ALK Protein in Different Cell Types

Two different approaches were used to elucidate which ALK's are physiological type I receptors for TGF- $\beta$ .

Firstly, several cell lines were tested for the expression of the ALK proteins by cross-linking followed by immunoprecipitation using the specific antisera against ALKs and the TGF- $\beta$  type II receptor. The mink lung epithelial cell line, Mv1Lu, is widely used to provide target cells for TGF- $\beta$  action and is well characterized regarding TGF- $\beta$  receptors (Laiho *et al* (1990) J. Biol. Chem. 265, 18518-18524; Laiho *et al* (1991) J. Biol. Chem. 266, 9108-9112). Only the VPN antiserum efficiently precipitated both type I and type II TGF- $\beta$  receptors in the wild type Mv1Lu cells. The DRL antiserum also precipitated components with the same size as those precipitated by the VPN antiserum. A mutant cell line (R mutant) which lacks the TGF- $\beta$  type I receptor and does not respond to TGF- $\beta$  (Laiho *et al*, *supra*) was also investigated by cross-linking followed by immunoprecipitation. Consistent with the results obtained by Laiho *et al* (1990), *supra* the type III and type II TGF- $\beta$  receptor complexes, but not the type I receptor complex, were observed by affinity crosslinking. Crosslinking followed by immunoprecipitation using the DRL antiserum revealed only the type II receptor complex, whereas neither the type I nor type II receptor complexes was seen using the VPN antiserum. When the cells were metabolically labelled and subjected to immunoprecipitation using the VPN antiserum, the 53 kDa ALK-5 protein was precipitated in both the wild-type and R mutant Mv1Lu cells. These results suggest that the type I receptor expressed in the R mutant is ALK-5, which has lost the affinity for binding to TGF- $\beta$  after mutation.



The type I and type II TGF- $\beta$  receptor complexes could be precipitated by the VPN and DRL antisera in other cell lines, including human foreskin fibroblasts (AG1518), human lung adenocarcinoma cells (A549), and human oral squamous cell carcinoma cells (HSC-2). Affinity cross-linking studies revealed multiple TGF- $\beta$  type I receptor-like complexes of 70-77 kDa in these cells. These components were less efficiently competed by excess unlabelled TGF- $\beta$ 1 in HSC-2 cells. Moreover, the type II receptor complex was low or not detectable in A549 and HSC-2 cells. Cross-linking followed by immunoprecipitation revealed that the VPN antiserum precipitated only the 70 kDa complex among the 70-77 kDa components. The DRL antiserum precipitated the 94 kDa type II receptor complex as well as the 70 kDa type I receptor complex in these cells, but not the putative type I receptor complexes of slightly larger sizes. These results suggest that multiple type I TGF- $\beta$  receptors may exist and that the 70 kDa complex containing ALK-5 forms a heteromeric complex with the TGF- $\beta$  type II receptor cloned by Lin *et al* (1992) Cell 68, 775-785, more efficiently than the other species. In rat pheochromocytoma cells (PC12) which have been reported to have no TGF- $\beta$  receptor complexes by affinity cross-linking (Massagué *et al* (1990) Ann. N.Y. Acad. Sci. 593, 59-72), neither VPN nor DRL antisera precipitated the TGF- $\beta$  receptor complexes. The antisera against ALKs -1 to -4 and ALK6 did not efficiently immunoprecipitate the crosslinked receptor complexes in porcine aortic endothelial (PAE) cells or human foreskin fibroblasts.

Next, it was investigated whether ALKs could restore responsiveness to TGF- $\beta$  in the R mutant of Mv1Lu cells, which lack the ligand-binding ability of the TGF- $\beta$  type I receptor but have intact type II receptor. Wild-type Mv1Lu cells and mutant cells were transfected with ALK cDNA and were then assayed for the production of plasminogen activator inhibitor-1 (PAI-1) which is produced as a result of TGF- $\beta$  receptor activation as described previously by

Laiho *et al* (1991) Mol. Cell Biol. 11, 972-978. Briefly, cells were added with or without 10 ng/ml of TGF- $\beta$ 1 for 2 hours in serum-free MCDB 104 without methionine. Thereafter, cultures were labelled with [ $^{35}$ S] methionine (40  $\mu$ Ci/ml) for 2 hours. The cells were removed by washing on ice once in PBS, twice in 10 mM Tris-HCl (pH 8.0), 0.5% sodium deoxycholate, 1 mM PMSF, twice in 2 mM Tris-HCl (pH 8.0), and once in PBS. Extracellular matrix proteins were extracted by scraping cells into the SDS-sample buffer containing DTT, and analyzed by SDS-gel electrophoresis followed by fluorography using Amplify. PAI-1 can be identified as a characteristic 45kDa band (Laiho *et al* (1991) Mol. Cell Biol. 11, 972-978). Wild-type Mv1Lu cells responded to TGF- $\beta$  and produced PAI-1, whereas the R mutant clone did not, even after stimulation by TGF- $\beta$ 1. Transient transfection of the ALK-5 cDNA into the R mutant clone led to the production of PAI-1 in response to the stimulation by TGF- $\beta$ 1, indicating that the ALK-5 cDNA encodes a functional TGF- $\beta$  type I receptor. In contrast, the R mutant cells that were transfected with other ALKs did not produce PAI-1 upon the addition of TGF- $\beta$ 1.

Using similar approaches as those described above for the identification of TGF- $\beta$ -binding ALKs, the ability of ALKs to bind activin in the presence of ActRII was examined. COS-1 cells were co-transfected as described above. Recombinant human activin A was iodinated using the chloramine T method (Mathews and Vale (1991) Cell 65, 973-982). Transfected COS-1 cells were analysed for binding and crosslinking of  $^{125}$ I-activin A in the presence or absence of excess unlabelled activin A. The crosslinked complexes were subjected to immunoprecipitation using DRL antisera or specific ALK antisera.

All ALKs appear to bind activin A in the presence of Act R-II. This is more clearly demonstrated by affinity cross-linking followed by immunoprecipitation. ALK-2 and ALK-4 bound  $^{125}$ I-activin A and were coimmunoprecipitated

with ActR-II. Other ALKs also bound <sup>125</sup>I-activin A but with a lower efficiency compared to ALK-2 and ALK-4.

In order to investigate whether ALKs are physiological activin type I receptors, activin responsive cells were examined for the expression of endogenous activin type I receptors. Mv1Lu cells, as well as the R mutant, express both type I and type II receptors for activin, and the R mutant cells produce PAI-1 upon the addition of activin A. Mv1Lu cells were labeled with <sup>125</sup>I-activin A, cross-linked and immunoprecipitated by the antisera against ActR-II or ALKs as described above.

The type I and type II receptor complexes in Mv1Lu cells were immunoprecipitated only by the antisera against ALK-2, ALK-4 and ActR-II. Similar results were obtained using the R mutant cells. PAE cells do not bind activin because of the lack of type II receptors for activin, and so cells were transfected with a chimeric receptor, to enable them to bind activin, as described herein. A plasmid (chim A) containing the extracellular domain and C-terminal tail of Act R-II (amino-acids -19 to 116 and 465 to 494, respectively (Mathews and Vale (1991) Cell, 65, 973-982)) and the kinase domain of TBR-II (amino-acids 160-543) (Lin et al (1992) Cell, 68, 775-785) was constructed and transfected into pCDNA/neo (Invitrogen). PAE cells were stably transfected with the chim A plasmid by electroporation, and cells expressing the chim A protein were established as described previously. PAE/Chim A cells were then subjected to <sup>125</sup>I-activin A labelling crosslinking and immunoprecipitation as described above.

Similar to Mv1Lu cells, activin type I receptor complexes in PAE/Chim A cells were immunoprecipitated by the ALK-2 and ALK-4 antisera. These results show that both ALK-2 and ALK-4 serve as high affinity type I receptors for activin A in these cells.

ALK-1, ALK-3 and ALK-6 bind TGF-β1 and activin A in the presence of their respective type II receptors, but the

functional consequences of the binding of the ligands remains to be elucidated.

The invention has been described by way of example only, without restriction of its scope. The invention is  
5 defined by the subject matter herein, including the claims that follow the immediately following full Sequence Listings:

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Ludwig Institute for Cancer Research
- (B) STREET: St. Mary's Hospital Medical School, Norfolk Place
- (C) CITY: Paddington, London
- (E) COUNTRY: United Kingdom
- (F) POSTAL CODE (ZIP): W2 1PG

(ii) TITLE OF INVENTION: PROTEINS HAVING SERINE/THREONINE KINASE DOMAINS, CORRESPONDING NUCLEIC ACID MOLECULES, AND THEIR USE

(iii) NUMBER OF SEQUENCES: 29

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1984 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 283..1791

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

AGGAAACGGT TTATTAGGAG GGAGTGGTGG AGCTGGGCCA GGCAGGAAGA CGCTGGAATA	60
AGAAACATTT TTGCTCCAGC CCCCATCCCA GTCCCGGGAG GCTGCCGCCC CAGCTGCGCC	120
GAGCGAGCCC CTCCCCGGCT CCAGCCCGGT CCGGGGCCGC GCCGGACCCC AGCCCGCCGT	180
CCAGCGCTGG CGGTGCAACT GCGGCGCGC GGTGGAGGGG AGGTGGCCCC GGTCCGCCGA	240



AAG ATC TTC TCC TCG AGG GAT GAA CAG TCC TGG TTC CGG GAG ACT GAG Lys Ile Phe Ser Ser Arg Asp Glu Gln Ser Trp Phe Arg Glu Thr Glu 230 235 240	1014
ATC TAT AAC ACA GTA TTG CTC AGA CAC GAC AAC ATC CTA GGC TTC ATC Ile Tyr Asn Thr Val Leu Leu Arg His Asp Asn Ile Leu Gly Phe Ile 245 250 255 260	1062
GCC TCA GAC ATG ACC TCC CGC AAC TCG AGC ACG CAG CTG TGG CTC ATC Ala Ser Asp Met Thr Ser Arg Asn Ser Ser Thr Gln Leu Trp Leu Ile 265 270 275	1110
ACG CAC TAC CAC GAG CAC GGC TCC CTC TAC GAC TTT CTG CAG AGA CAG Thr His Tyr-His Glu His-Gly-Ser Leu Tyr Asp Phe Leu Gln Arg Gln 280 285 290	1158
ACG CTG GAG CCC CAT CTG GCT CTG AGG CTA GCT GTG TCC GCG GCA TGC Thr Leu Glu Pro His Leu Ala Leu Arg Leu Ala Val Ser Ala Ala Cys 295 300 305	1206
GGC CTG GCG CAC CTG CAC GTG GAG ATC TTC GGT ACA CAG GGC AAA CCA Gly Leu Ala His Leu His Val Glu Ile Phe Gly Thr Gln Gly Lys Pro 310 315 320	1254
GCC ATT GCC CAC CGC GAC TTC AAG AGC CGC AAT GTG CTG GTC AAG AGC Ala Ile Ala His Arg Asp Phe Lys Ser Arg Asn Val Leu Val Lys Ser 325 330 335 340	1302
AAC CTG CAG TGT TGC ATC GCC GAC CTG GGC CTG GCT GTG ATG CAC TCA Asn Leu Gln Cys Cys Ile Ala Asp Leu Gly Leu Ala Val Met His Ser 345 350 355	1350
CAG GGC AGC GAT TAC CTG GAC ATC GGC AAC AAC CCG AGA GTG GGC ACC Gln-Gly-Ser Asp Tyr Leu Asp Ile Gly Asn Asn Pro Arg Val Gly Thr 360 365 370	1398
AAG CGG TAC ATG GCA CCC GAG GTG CTG GAC GAG CAG ATC CGC ACG GAC Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Gln Ile Arg Thr Asp 375 380 385	1446
TGC TTT GAG TCC TAC AAG TGG ACT GAC ATC TGG GCC TTT GGC CTG GTG Cys Phe Glu Ser Tyr Lys Trp Thr Asp Ile Trp Ala Phe Gly Leu Val 390 395 400	1494
CTG TGG GAG ATT GCC CGC CGG ACC ATC GTG AAT GGC ATC GTG GAG GAC Leu Trp Glu Ile Ala Arg Arg Thr Ile Val Asn Gly Ile Val Glu Asp 405 410 415 420	1542
TAT AGA CCA CCC TTC TAT GAT GTG GTG CCC AAT GAC CCC AGC TTT GAG Tyr Arg Pro Pro Phe Tyr Asp Val Val Pro Asn Asp Pro Ser Phe Glu 425 430 435	1590
GAC ATG AAG AAG GTG GTG TGT GTG GAT CAG CAG ACC CCC ACC ATC CCT Asp Met Lys Lys Val Val Cys Val Asp Gln Gln Thr Pro Thr Ile Pro 440 445 450	1638
AAC CGG CTG GCT GCA GAC CCG GTC CTC TCA GGC CTA GCT CAG ATG ATG Asn Arg Leu Ala Ala Asp Pro Val Leu Ser Gly Leu Ala Gln Met Met 455 460 465	1686

38

CGG GAG TGC TGG TAC CCA AAC CCC TCT GCC CGA CTC ACC GCG CTG CGG 1734  
 Arg Glu Cys Trp Tyr Pro Asn Pro Ser Ala Arg Leu Thr Ala Leu Arg  
 470 475 480  
 ATC AAG AAG ACA CTA CAA AAA ATT AGC AAC AGT CCA GAG AAG CCT AAA 1782  
 Ile Lys Lys Thr Leu Gln Lys Ile Ser Asn Ser Pro Glu Lys Pro Lys  
 485 490 495 500  
 GTG ATT CAA TAGCCCAGGA GCACCTGATT CCTTTCTGCC TGCAGGGGGC 1831  
 Val Ile Gln  
 TGGGGGGGTG GGGGGCAGTG GATGGTGCCC TATCTGGGTA GAGGTAGTGT GAGTGTGGTG 1891  
 TGTGCTGGGG ATGGGCAGCT GCGCCTGCCT GCTCGGCCCC CAGCCCACCC AGCCAAAAAT 1951  
 ACAGCTGGGC TGAAACCTGA AAAAAAAAAA AAA 1984

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 503 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Thr Leu Gly Ser Pro Arg Lys Gly Leu Leu Met Leu Leu Met Ala  
 1 5 10 15  
 Leu Val Thr Gln Gly Asp Pro Val Lys Pro Ser Arg Gly Pro Leu Val  
 20 25 30  
 Thr Cys Thr Cys Glu Ser Pro His Cys Lys Gly Pro Thr Cys Arg Gly  
 35 40 45  
 Ala Trp Cys Thr Val Val Leu Val Arg Glu Glu Gly Arg His Pro Gln  
 50 55 60  
 Glu His Arg Gly Cys Gly Asn Leu His Arg Glu Leu Cys Arg Gly Arg  
 65 70 75 80  
 Pro Thr Glu Phe Val Asn His Tyr Cys Cys Asp Ser His Leu Cys Asn  
 85 90 95  
 His Asn Val Ser Leu Val Leu Glu Ala Thr Gln Pro Pro Ser Glu Gln  
 100 105 110  
 Pro Gly Thr Asp Gly Gln Leu Ala Leu Ile Leu Gly Pro Val Leu Ala  
 115 120 125  
 Leu Leu Ala Leu Val Ala Leu Gly Val Leu Gly Leu Trp His Val Arg  
 130 135 140  
 Arg Arg Gln Glu Lys Gln Arg Gly Leu His Ser Glu Leu Gly Glu Ser  
 145 150 155 160

SUBSTITUTE SHEET



Ser Leu Ile Leu Lys Ala Ser Glu Gln Gly Asp Thr Met Leu Gly Asp  
 165 170 175  
 Leu Leu Asp Ser Asp Cys Thr Thr Gly Ser Gly Ser Gly Leu Pro Phe  
 180 185 190  
 Leu Val Gln Arg Thr Val Ala Arg Gln Val Ala Leu Val Glu Cys Val  
 195 200 205  
 Gly Lys Gly Arg Tyr Gly Glu Val Trp Arg Gly Leu Trp His Gly Glu  
 210 215 220  
 Ser Val Ala Val Lys Ile Phe Ser Ser Arg Asp Glu Gln Ser Trp Phe  
 225 230 235 240  
 Arg Glu Thr Glu Ile Tyr Asn Thr Val Leu Leu Arg His Asp Asn Ile  
 245 250 255  
 Leu Gly Phe Ile Ala Ser Asp Met Thr Ser Arg Asn Ser Ser Thr Gln  
 260 265 270  
 Leu Trp Leu Ile Thr His Tyr His Glu His Gly Ser Leu Tyr Asp Phe  
 275 280 285  
 Leu Gln Arg Gln Thr Leu Glu Pro His Leu Ala Leu Arg Leu Ala Val  
 290 295 300  
 Ser Ala Ala Cys Gly Leu Ala His Leu His Val Glu Ile Phe Gly Thr  
 305 310 315 320  
 Gln Gly Lys Pro Ala Ile Ala His Arg Asp Phe Lys Ser Arg Asn Val  
 325 330 335  
 Leu Val Lys Ser Asn Leu Gln Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 340 345 350  
 Val Met His Ser Gln Gly Ser Asp Tyr Leu Asp Ile Gly Asn Asn Pro  
 355 360 365  
 Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Gln  
 370 375 380  
 Ile Arg Thr Asp Cys Phe Glu Ser Tyr Lys Trp Thr Asp Ile Trp Ala  
 385 390 395 400  
 Phe Gly Leu Val Leu Trp Glu Ile Ala Arg Arg Thr Ile Val Asn Gly  
 405 410 415  
 Ile Val Glu Asp Tyr Arg Pro Pro Phe Tyr Asp Val Val Pro Asn Asp  
 420 425 430  
 Pro Ser Phe Glu Asp Met Lys Lys Val Val Cys Val Asp Gln Gln Thr  
 435 440 445  
 Pro Thr Ile Pro Asn Arg Leu Ala Ala Asp Pro Val Leu Ser Gly Leu  
 450 455 460  
 Ala Gln Met Met Arg Glu Cys Trp Tyr Pro Asn Pro Ser Ala Arg Leu  
 465 470 475 480

40

Thr Ala Leu Arg Ile Lys Lys Thr Leu Gln Lys Ile Ser Asn Ser Pro  
 485 490 495

Glu Lys Pro Lys Val Ile Gln  
 500

## (2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2724 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: Homo sapiens

(ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 104..1630

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CTCCGAGTAC CCCAGTGACC AGAGTGAGAG AAGCTCTGAA CGAGGGCACC CGGCTTGAAG	60
GA CTGTGGGC AGATGTGACC AAGAGCCTGC ATTAAGTTGT ACA ATG GTA GAT GGA	115
Met Val Asp Gly	
1	
GTG ATG ATT CTT CCT GTG CTT ATC ATG ATT GCT CTC CCC TCC CCT AGT	163
Val Met Ile Leu Pro Val Leu Ile Met Ile Ala Leu Pro Ser Pro Ser	
5 10 15 20	
ATG GAA GAT GAG AAG CCC AAG GTC AAC CCC AAA CTC TAC ATG TGT GTG	211
Met Glu Asp Glu Lys Pro Lys Val Asn Pro Lys Leu Tyr Met Cys Val	
25 30 35	
TGT GAA GGT CTC TCC TGC GGT AAT GAG GAC CAC TGT GAA GGC CAG CAG	259
Cys Glu Gly Leu Ser Cys Gly Asn Glu Asp His Cys Glu Gly Gln Gln	
40 45 50	
TGC TTT TCC TCA CTG AGC ATC AAC GAT GGC TTC CAC GTC TAC CAG AAA	307
Cys Phe Ser Ser Leu Ser Ile Asn Asp Gly Phe His Val Tyr Gln Lys	
55 60 65	
GGC TGC TTC CAG GTT TAT GAG CAG GGA AAG ATG ACC TGT AAG ACC CCG	355
Gly Cys Phe Gln Val Tyr Glu Gln Gly Lys Met Thr Cys Lys Thr Pro	
70 75 80	

CCG Pro 85	TCC Ser	CCT Pro	GGC Gly	CAA Gln	GCT Ala	GTG Val	GAG Glu	TGC Cys	TGC Cys	CAA Gln	GGG Gly	GAC Asp	TGG Trp	TGT Cys	AAC Asn	403
					90					95					100	
AGG Arg	AAC Asn	ATC Ile	ACG Thr	GCC Ala	CAG Gln	CTG Leu	CCC Pro	ACT Thr	AAA Lys	GGA Gly	AAA Lys	TCC Ser	TTC Phe	CCT Pro	GGA Gly	451
				105					110					115		
ACA Thr	CAG Gln	AAT Asn	TTC Phe	CAC His	TTG Leu	GAG Glu	GTT Val	GGC Gly	CTC Leu	ATT Ile	ATT Ile	CTC Leu	TCT Ser	GTA Val	GTG Val	499
			120					125					130			
TTC Phe	GCA Ala	GTA Val	TGT Cys	CTT Leu	TTA Leu	GCC Ala	TGC Cys	CTG Leu	CTG Leu	GGA Gly	GTT Val	GCT Ala	CTC Leu	CGA Arg	AAA Lys	547
		135					140					145				
TTT Phe	AAA Lys	AGG Arg	CGC Arg	AAC Asn	CAA Gln	GAA Glu	CGC Arg	CTC Leu	AAT Asn	CCC Pro	CGA Arg	GAC Asp	GTG Val	GAG Glu	TAT Tyr	595
	150					155					160					
GGC Gly	ACT Thr	ATC Ile	GAA Glu	GGG Gly	CTC Leu	ATC Ile	ACC Thr	ACC Thr	AAT Asn	GTT Val	GGA Gly	GAC Asp	AGC Ser	ACT Thr	TTA Leu	643
165					170					175					180	
GCA Ala	GAT Asp	TTA Leu	TTG Leu	GAT Asp	CAT His	TCG Ser	TGT Cys	ACA Thr	TCA Ser	GGA Gly	AGT Ser	GGC Gly	TCT Ser	GGT Gly	CTT Leu	691
				185					190					195		
CCT Pro	TTT Phe	CTG Leu	GTA Val	CAA Gln	AGA Arg	ACA Thr	GTG Val	GCT Ala	CGC Arg	CAG Gln	ATT Ile	ACA Thr	CTG Leu	TTG Leu	GAG Glu	739
			200					205					210			
TGT Cys	GTC Val	GGG Gly	AAA Lys	GGC Gly	AGG Arg	TAT Tyr	GGT Gly	GAG Glu	GTG Val	TGG Trp	AGG Arg	GGC Gly	AGC Ser	TGG Trp	CAA Gln	787
		215					220					225				
GGG Gly	GAA Glu	AAT Asn	GTT Val	GCC Ala	GTG Val	AAG Lys	ATC Ile	TTC Phe	TCC Ser	TCC Ser	CGT Arg	GAT Asp	GAG Glu	AAG Lys	TCA Ser	835
	230					235					240					
TGG Trp	TTC Phe	AGG Arg	GAA Glu	ACG Thr	GAA Glu	TTG Leu	TAC Tyr	AAC Asn	ACT Thr	GTG Val	ATG Met	CTG Leu	AGG Arg	CAT His	GAA Glu	883
245					250					255					260	
AAT Asn	ATC Ile	TTA Leu	GGT Gly	TTC Phe	ATT Ile	GCT Ala	TCA Ser	GAC Asp	ATG Met	ACA Thr	TCA Ser	AGA Arg	CAC His	TCC Ser	AGT Ser	931
				265					270					275		
ACC Thr	CAG Gln	CTG Leu	TGG Trp	TTA Leu	ATT Ile	ACA Thr	CAT His	TAT Tyr	CAT His	GAA Glu	ATG Met	GGA Gly	TCG Ser	TTG Leu	TAC Tyr	979
			280					285					290			
GAC Asp	TAT Tyr	CTT Leu	CAG Gln	CTT Leu	ACT Thr	ACT Thr	CTG Leu	GAT Asp	ACA Thr	GTT Val	AGC Ser	TGC Cys	CTT Leu	CGA Arg	ATA Ile	1027
		295					300					305				
GTG Val	CTG Leu	TCC Ser	ATA Ile	GCT Ala	AGT Ser	GGT Gly	CTT Leu	GCA Ala	CAT His	TTG Leu	CAC His	ATA Ile	GAG Glu	ATA Ile	TTT Phe	1075
	310					315					320					

GGG ACC CAA GGG AAA CCA GCC ATT GCC CAT CGA GAT TTA AAG AGC AAA Gly Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys 325 330 335 340	1123
AAT ATT CTG GTT AAG AAG AAT GGA CAG TGT TGC ATA GCA GAT TTG GGC Asn Ile Leu Val Lys Lys Asn Gly Gln Cys Cys Ile Ala Asp Leu Gly 345 350 355	1171
CTG GCA GTC ATG CAT TCC CAG AGC ACC AAT CAG CTT GAT GTG GGG AAC Leu Ala Val Met His Ser Gln Ser Thr Asn Gln Leu Asp Val Gly Asn 360 365 370	1219
AAT CCC CGT GTG GGC ACC AAG CGC TAC ATG GCC CCC GAA GTT CTA GAT Asn Pro Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp 375 380 385	1267
GAA ACC ATC CAG GTG GAT TGT TTC GAT TCT TAT AAA AGG GTC GAT ATT Glu Thr Ile Gln Val Asp Cys Phe Asp Ser Tyr Lys Arg Val Asp Ile 390 395 400	1315
TGG GCC TTT GGA CTT GTT TTG TGG GAA GTG GCC AGG CCG ATG GTG AGC Trp Ala Phe Gly Leu Val Leu Trp Glu Val Ala Arg Arg Met Val Ser 405 410 415 420	1363
AAT GGT ATA GTG GAG GAT TAC AAG CCA CCG TTC TAC GAT GTG GTT CCC Asn Gly Ile Val Glu Asp Tyr Lys Pro Pro Phe Tyr Asp Val Val Pro 425 430 435	1411
AAT GAC CCA AGT TTT GAA GAT ATG AGG AAG GTA GTC TGT GTG GAT CAA Asn Asp Pro Ser Phe Glu Asp Met Arg Lys Val Val Cys Val Asp Gln 440 445 450	1459
CAA AGG CCA AAC ATA CCC AAC AGA TGG TTC TCA GAC CCG ACA TTA ACC Gln Arg Pro Asn Ile Pro Asn Arg Trp Phe Ser Asp Pro Thr Leu Thr 455 460 465	1507
TCT CTG GCC AAG CTA ATG AAA GAA TGC TGG TAT CAA AAT CCA TCC GCA Ser Leu Ala Lys Leu Met Lys Glu Cys Trp Tyr Gln Asn Pro Ser Ala 470 475 480	1555
AGA CTC ACA GCA CTG CGT ATC AAA AAG ACT TTG ACC AAA ATT GAT AAT Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Thr Lys Ile Asp Asn 485 490 495 500	1603
TCC CTC GAC AAA TTG AAA ACT GAC TGT TGACATTTTC ATAGTGTCAA Ser Leu Asp Lys Leu Lys Thr Asp Cys 505	1650
GAAGGAAGAT TTGACGTTGT TGTCATTGTC CAGCTGGGAC CTAATGCTGG CCTGACTGGT	1710
TGTCAGAATG GAATCCATCT GTCTCCCTCC CCAAATGGCT GCTTTGACAA GGCAGACGTC	1770
GTACCCAGCC ATGTGTTGGG GAGACATCAA AACCACCCTA ACCTCGCTCG ATGACTGTGA	1830
ACTGGGCATT TCACGAACTG TTCACACTGC AGAGACTAAT GTTGGACAGA CACTGTTGCA	1890
AAGGTAGGGA CTGGAGGAAC ACAGAGAAAT CCTAAAAGAG ATCTGGGCAT TAAGTCAGTG	1950
GCTTTGCATA GCTTTCACAA GTCTCCTAGA CACTCCCCAC GGGAAACTCA AGGAGGTGGT	2010

```

GAATTTTAA TCAGCAATAT TGCCTGTGCT TCTCTTCTTT ATTGCACTAG GAATTCTTTG      2070
CATTCCTTAC TTGCACTGTT ACTCTTAATT TTAAAGACCC AACTTGCCAA AATGTTGGCT      2130
GCGTACTCCA CTGGTCTGTC TTTGGATAAT AGGAATTCAA TTTGGCAAAA CAAAATGTAA      2190
TGTCAGACTT TGCTGCATTT TACACATGTG CTGATGTTTA CAATGATGCC GAACATTAGG      2250
AATTGTTTAT ACACAACCTT GCAAATTATT TATTACTTGT GCACTTAGTA GTTTTACAA      2310
AACTGCTTTG TGCATATGTT AAAGCTTATT TTTATGTGGT CTTATGATTT TATTACAGAA      2370
ATGTTTTTAA CACTATACTC TAAAATGGAC ATTTTCTTTT ATTATCAGTT AAAATCACAT      2430
TTTAAGTGCT TCACATTTGT ATGTGTGTAG ACTGTAACCT TTTTTCAGTT CATATGCAGA      2490
ACGTATTTAG CCATTACCCA CGTGACACCA CCGAATATAT TATCGATTTA GAAGCAAAGA      2550
TTTCAGTAGA ATTTTAGTCC TGAACGCTAC GGGGAAAATG CATTTTCTTC AGAATTATCC      2610
ATTACGTGCA TTAAACTCT GCCAGAAAAA AATAACTATT TTGTTTTAAT CTACTTTTGT      2670
TATTTAGTAG TTATTTGTAT AAATTAAATA AACTGTTTTC AAGTCAAPAA AAAA          2724

```

## (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 509 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Val Asp Gly Val Met Ile Leu Pro Val Leu Ile Met Ile Ala Leu
 1           5           10
Pro Ser Pro Ser Met Glu Asp Glu Lys Pro Lys Val Asn Pro Lys Leu
 20          25          30
Tyr Met Cys Val Cys Glu Gly Leu Ser Cys Gly Asn Glu Asp His Cys
 35          40          45
Glu Gly Gln Gln Cys Phe Ser Ser Leu Ser Ile Asn Asp Gly Phe His
 50          55          60
Val Tyr Gln Lys Gly Cys Phe Gln Val Tyr Glu Gln Gly Lys Met Thr
 65          70          75          80
Cys Lys Thr Pro Pro Ser Pro Gly Gln Ala Val Glu Cys Cys Gln Gly
 85          90          95
Asp Trp Cys Asn Arg Asn Ile Thr Ala Gln Leu Pro Thr Lys Gly Lys
100         105         110
Ser Phe Pro Gly Thr Gln Asn Phe His Leu Glu Val Gly Leu Ile Ile
115         120         125

```

Leu Ser Val Val Phe Ala Val Cys Leu Leu Ala Cys Leu Leu Gly Val  
 130 135 140  
 Ala Leu Arg Lys Phe Lys Arg Arg Asn Gln Glu Arg Leu Asn Pro Arg  
 145 150 155 160  
 Asp Val Glu Tyr Gly Thr Ile Glu Gly Leu Ile Thr Thr Asn Val Gly  
 165 170 175  
 Asp Ser Thr Leu Ala Asp Leu Leu Asp His Ser Cys Thr Ser Gly Ser  
 180 185 190  
 Gly Ser Gly Leu Pro Phe Leu Val Gln Arg Thr Val Ala Arg Gln Ile  
 195 200 205  
 Thr Leu Leu Glu Cys Val Gly Lys Gly Arg Tyr Gly Glu Val Trp Arg  
 210 215 220  
 Gly Ser Trp Gln Gly Glu Asn Val Ala Val Lys Ile Phe Ser Ser Arg  
 225 230 235 240  
 Asp Glu Lys Ser Trp Phe Arg Glu Thr Glu Leu Tyr Asn Thr Val Met  
 245 250 255  
 Leu Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ser Asp Met Thr Ser  
 260 265 270  
 Arg His Ser Ser Thr Gln Leu Trp Leu Ile Thr His Tyr His Glu Met  
 275 280 285  
 Gly Ser Leu Tyr Asp Tyr Leu Gln Leu Thr Thr Leu Asp Thr Val Ser  
 290 295 300  
 Cys Leu Arg Ile Val Leu Ser Ile Ala Ser Gly Leu Ala His Leu His  
 305 310 315 320  
 Ile Glu Ile Phe Gly Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp  
 325 330 335  
 Leu Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Gln Cys Cys Ile  
 340 345 350  
 Ala Asp Leu Gly Leu Ala Val Met His Ser Gln Ser Thr Asn Gln Leu  
 355 360 365  
 Asp Val Gly Asn Asn Pro Arg Val Gly Thr Lys Arg Tyr Met Ala Pro  
 370 375 380  
 Glu Val Leu Asp Glu Thr Ile Gln Val Asp Cys Phe Asp Ser Tyr Lys  
 385 390 395 400  
 Arg Val Asp Ile Trp Ala Phe Gly Leu Val Leu Trp Glu Val Ala Arg  
 405 410 415  
 Arg Met Val Ser Asn Gly Ile Val Glu Asp Tyr Lys Pro Pro Phe Tyr  
 420 425 430  
 Asp Val Val Pro Asn Asp Pro Ser Phe Glu Asp Met Arg Lys Val Val  
 435 440 445

45

Cys Val Asp Gln Gln Arg Pro Asn Ile Pro Asn Arg Trp Phe Ser Asp  
 450 455 460

Pro Thr Leu Thr Ser Leu Ala Lys Leu Met Lys Glu Cys Trp Tyr Gln  
 465 470 475 480

Asn Pro Ser Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Thr  
 485 490 495

Lys Ile Asp Asn Ser Leu Asp Lys Leu Lys Thr Asp Cys  
 500 505

## (2) INFORMATION FOR SEQ ID NO: 5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2932 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iii) ANTI-SENSE: NO

## (v) FRAGMENT TYPE: internal

## (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 310..1905

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

GCTCCGCGCC GAGGGCTGGA GGATGCGTTC CCTGGGGTCC GGACTTATGA AAATATGCAT	60
CAGTTTAATA CTGTCTTGGA ATTCATGAGA TGGAGGCATA GGTCAAAGCT GTTTGGAGAA	120
AATCAGAAGT ACAGTTTTAT CTAGCCACAT CTTGGAGGAG TCGTAAGAAA GCAGTGGGAG	180
TTGAAGTCAT TGTCAAGTGC TTGCGATCTT TTACAAGAAA ATCTCACTGA ATGATAGTCA	240
TTTAAATTGG TGAAGTAGCA AGACCAATTA TTAAGGTGA CAGTACACAG GAAACATTAC	300
AATTGAACA ATG ACT CAG CTA TAC ATT TAC ATC AGA TTA TTG GGA GCC	348
Met Thr Gln Leu Tyr Ile Tyr Ile Arg Leu Leu Gly Ala	
1 5 10	
TAT TTG TTC ATC ATT TCT CGT GTT CAA GGA CAG AAT CTG GAT AGT ATG	396
Tyr Leu Phe Ile Ile Ser Arg Val Gln Gly Gln Asn Leu Asp Ser Met	
15 20 25	

CTT CAT GGC ACT GGG ATG AAA TCA CAC TCC GAC CAG AAA AAG TCA GAA Leu His Gly Thr Gly Met Lys Ser Asp Ser Asp Gln Lys Lys Ser Glu 30 35 40 45	444
AAT GGA GTA ACC TTA GCA CCA GAG GAT ACC TTG CCT TTT TTA AAG TGC Asn Gly Val Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys 50 55 60	492
TAT TGC TCA GGG CAC TGT CCA GAT GAT GCT ATT AAT AAC ACA TGC ATA Tyr Cys Ser Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile 65 70 75	540
ACT AAT GGA CAT TGC TTT GCC ATC ATA GAA GAA GAT GAC CAG GGA GAA Thr Asn Gly His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu 80 85 90	588
ACC ACA TTA GCT TCA GGG TGT ATG AAA TAT GAA GGA TCT GAT TTT CAG Thr Thr Leu Ala Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln 95 100 105	636
TGC AAA GAT TCT CCA AAA GCC CAG CTA CGC CGG ACA ATA GAA TGT TGT Cys Lys Asp Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys 110 115 120 125	684
CGG ACC AAT TTA TGT AAC CAG TAT TTG CAA CCC ACA CTG CCC CCT GTT Arg Thr Asn Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val 130 135 140	732
GTC ATA GGT CCG TTT TTT GAT GGC AGC ATT CGA TGG CTG GTT TTG CTC Val Ile Gly Pro Phe Phe Asp Gly Ser Ile Arg Trp Leu Val Leu Leu 145 150 155	780
ATT TCT ATG GCT GTC TGC ATA ATT GCT ATG ATC ATC TTC TCC AGC TGC Ile Ser Met Ala Val Cys Ile Ile Ala Met Ile Ile Phe Ser Ser Cys 160 165 170	828
TTT TGT TAC AAA CAT TAT TGC AAG AGC ATC TCA AGC AGA CGT CGT TAC Phe Cys Tyr Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Arg Arg Tyr 175 180 185	876
AAT CGT GAT TTG GAA CAG GAT GAA GCA TTT ATT CCA GTT GGA GAA TCA Asn Arg Asp Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser 190 195 200 205	924
CTA AAA GAC CTT ATT GAC CAG TCA CAA AGT TCT GGT AGT GGG TCT GGA Leu Lys Asp Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly 210 215 220	972
CTA CCT TTA TTG GTT CAG CGA ACT ATT GCC AAA CAG ATT CAG ATG GTC Leu Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val 225 230 235	1020
CGG CAA GTT GGT AAA GGC CGA TAT GGA GAA GTA TGG ATG GGC AAA TGG Arg Gln Val Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp 240 245 250	1068
CGT GGC GAA AAA GTG GCG GTG AAA GTA TTC TTT ACC ACT GAA GAA GCC Arg Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala 255 260 265	1116



AGC Ser 270	TGG Trp	TTT Phe	CGA Arg	GAA Glu	ACA Thr	GAA Glu	ATC Ile	TAC Tyr	CAA Gln	ACT Thr	GTG Val	CTA Leu	ATG Met	CGC Arg	CAT His 285	1164
GAA Glu	AAC Asn	ATA Ile	CTT Leu	GGT Gly 290	TTC Phe	ATA Ile	GCG Ala	GCA Ala	GAC Asp 295	ATT Ile	AAA Lys	GGT Gly	ACA Thr	GGT Gly	TCC Ser 300	1212
TGG Trp	ACT Thr	CAG Gln	CTC Leu	TAT Tyr	TTG Leu	ATT Ile	ACT Thr	GAT Asp	TAC Tyr	CAT His	GAA Glu	AAT Asn	GGA Gly	TCT Ser	CTC Leu	1260
TAT Tyr	GAC Asp	TTC Phe	CTG Leu	AAA Lys	TGT Cys	GCT Ala	ACA Thr	CTG Leu	GAC Asp	ACC Thr	AGA Arg	GCC Ala	CTG Leu	CTT Leu	AAA Lys	1308
TTG Leu	GCT Ala	TAT Tyr	TCA Ser	GCT Ala	GCC Ala	TGT Cys	GGT Gly	CTG Leu	TGC Cys	CAC His	CTG Leu	CAC His	ACA Thr	GAA Glu	ATT Ile	1356
TAT Tyr	GGC Gly	ACC Thr	CAA Gln	GGA Gly	AAG Lys	CCC Pro	GCA Ala	ATT Ile	GCT Ala	CAT His	CGA Arg	GAC Asp	CTA Leu	AAG Lys	AGC Ser 365	1404
AAA Lys	AAC Asn	ATC Ile	CTC Leu	ATC Ile	AAG Lys	AAA Lys	AAT Asn	GGG Gly	AGT Ser	TGC Cys	TGC Cys	ATT Ile	GCT Ala	GAC Asp	CTG Leu	1452
GGC Gly	CTT Leu	GCT Ala	GTT Val	AAA Lys	TTC Phe	AAC Asn	AGT Ser	GAC Asp	ACA Thr	AAT Asn	GAA Glu	GTT Val	GAT Asp	GTG Val	CCC Pro	1500
TTG Leu	AAT Asn	ACC Thr	AGG Arg	GTG Val	GGC Gly	ACC Thr	AAA Lys	CGC Arg	TAC Tyr	ATG Met	GCT Ala	CCC Pro	GAA Glu	GTG Val	CTG Leu	1548
GAC Asp	GAA Glu	AGC Ser	CTG Leu	AAC Asn	AAA Lys	AAC Asn	CAC His	TTC Phe	CAG Gln	CCC Pro	TAC Tyr	ATC Ile	ATG Met	GCT Ala	GAC Asp	1596
ATC Ile	TAC Tyr	AGC Ser	TTC Phe	GGC Gly	CTA Leu	ATC Ile	ATT Ile	TGG Trp	GAG Glu	ATG Met	GCT Ala	CGT Arg	CGT Arg	TGT Cys	ATC Ile	1644
ACA Thr	GGA Gly	GGG Gly	ATC Ile	GTG Val	GAA Glu	GAA Glu	TAC Tyr	CAA Gln	TTG Leu	CCA Pro	TAT Tyr	TAC Tyr	AAC Asn	ATG Met	GTA Val	1692
CCG Pro	AGT Ser	GAT Asp	CCG Pro	TCA Ser	TAC Tyr	GAA Glu	GAT Asp	ATG Met	CGT Arg	GAG Glu	GTT Val	GTG Val	TGT Cys	GTC Val	AAA Lys	1740
CGT Arg	TTG Leu	CGG Arg	CCA Pro	ATT Ile	GTG Val	TCT Ser	AAT Asn	CGG Arg	TGG Trp	AAC Asn	AGT Ser	GAT Asp	GAA Glu	TGT Cys	CTA Leu	1788
CGA Arg	GCA Ala	GTT Val	TTG Leu	AAG Lys	CTA Leu	ATG Met	TCA Ser	GAA Glu	TGC Cys	TGG Trp	GCC Ala	CAC His	AAT Asn	CCA Pro	GCC Ala	1836

TCC AGA CTC ACA GCA TTG AGA ATT AAG AAG ACG CTT GCC AAG ATG GTT	1884
Ser Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val	
510 515 520 525	
GAA TCC CAA GAT GTA AAT ATC TGATGGTTAA ACCATCGGAG GAGAAACTCT	1935
Glu Ser Gln Asp Val Lys Ile	
530	
AGACTGCAAG AACTGTTTTT ACCCATGGCA TGGGTGGAAT TAGAGTGGAA TAAGGATGTT	1995
AACTTGGTTC TCAGACTCTT TCTTCACTAC GTGTTCCACAG GCTGCTAATA TTAAACCTTT	2055
CAGTACTCTT ATTAGGATAC AAGCTGGGAA CTTCTAAACA CTTTATTCTT TATATATGGA	2115
CAGCTTTATT TTAAATGTGG TTTTGTATGC CTTTTTTTAA GTGGGTTTTT ATGAACTGCA	2175
TCAAGACTTC AATCCTGATT AGTGTCTCCA GTCAAGCTCT GGGTACTGAA TTGCCTGTTC	2235
ATAAAACGGT GCTTCTGTG AAAGCCTTAA GAAGATAAAT GAGCGCAGCA GAGATGGAGA	2295
AATAGACTTT GCCTTTTACC TGAGACATTC AGTTCGTTTG TATTCTACCT TTGTAAAACA	2355
GCCTATAGAT GATGATGTGT TTGGGATACT GCTTATTTTA TGATAGTTTG TCCTGTGTCC	2415
TTAGTGATGT GTGTGTGTCT CCATGCACAT GCACGCCGGG ATTCCTCTGC TGCCATTGTA	2475
ATTAGAAGAA AATAATTTAT ATGCATGCAC AGGAAGATAT TGGTGGCCGG TGGTTTTGTG	2535
CTTTAAAAAT GCAATATCTG ACCAAGATTC GCCAATCTCA TACAAGCCAT TTACTTTGCA	2595
AGTGAGATAG CTTCCCCACC AGCTTTATTT TTTAACATGA AAGCTGATGC CAAGGCCAAA	2655
AGAAGTTTAA AGCATCTGTA AATTTGGACT GTTTTCCTTC AACCACCATT TTTTTGTGG	2715
TTATTATTTT TGTCACGGAA AGCATCCTCT CCAAAGTTGG AGCTTCTATT GCCATGAACC	2775
ATGCTTACAA AGAAAGCACT TCTTATTGAA GTGAATTCCT GCATTTGATA GCAATGTAAG	2835
TGCCTATAAC CATGTTCTAT ATTCTTTATT CTCAGTAACT TTTAAAAGGG AAGTTATTTA	2895
TATTTTGTGT ATAATGTGCT TTATTGCAA ATCACCC	2932

## (2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 532 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Met	Thr	Gln	Leu	Tyr	Ile	Tyr	Ile	Arg	Leu	Leu	Gly	Ala	Tyr	Leu	Phe
1				5				10						15	
Ile	Ile	Ser	Arg	Val	Gln	Gly	Gln	Asn	Leu	Asp	Ser	Met	Leu	His	Gly
			20					25					30		

Thr Gly Met Lys Ser Asp Ser Asp Gln Lys Lys Ser Glu Asn Gly Val  
 35 40 45  
 Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser  
 50 55 60  
 Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly  
 65 70 75 80  
 His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu  
 85 90 95  
 Ala Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp  
 100 105 110  
 Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn  
 115 120 125  
 Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly  
 130 135 140  
 Pro Phe Phe Asp Gly Ser Ile Arg Trp Leu Val Leu Leu Ile Ser Met  
 145 150 155 160  
 Ala Val Cys Ile Ile Ala Met Ile Ile Phe Ser Ser Cys Phe Cys Tyr  
 165 170 175  
 Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Arg Arg Tyr Asn Arg Asp  
 180 185 190  
 Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp  
 195 200 205  
 Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu  
 210 215 220  
 Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val  
 225 230 235 240  
 Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu  
 245 250 255  
 Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
 260 265 270  
 Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile  
 275 280 285  
 Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln  
 290 295 300  
 Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe  
 305 310 315 320  
 Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr  
 325 330 335  
 Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr  
 340 345 350

Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile  
 355 360 365  
 Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 370 375 380  
 Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Val Pro Leu Asn Thr  
 385 390 395 400  
 Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser  
 405 410 415  
 Leu Asn Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
 420 425 430  
 Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly  
 435 440 445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
 450 455 460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
 465 470 475 480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
 485 490 495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
 500 505 510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
 515 520 525  
 Asp Val Lys Ile  
 530

---

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2333 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1515

## (x1) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

ATG GCG GAG TCG GCC GGA GCC TCC TCC TTC TTC CCC CTT GTT GTC CTC	48
Met Ala Glu Ser Ala Gly Ala Ser Ser Phe Phe Pro Leu Val Val Leu	
1 5 10 15	
CTG CTC GCC GGC AGC GGC GGG TCC GGG CCC CGG GGG GTC CAG GCT CTG	96
Leu Leu Ala Gly Ser Gly Gly Ser Gly Pro Arg Gly Val Gln Ala Leu	
20 25 30	
CTG TGT GCG TGC ACC AGC TGC CTC CAG GCC AAC TAC ACG TGT GAG ACA	144
Leu Cys Ala Cys Thr Ser Cys Leu Gln Ala Asn Tyr Thr Cys Glu Thr	
35 40 45	
GAT GGG GCC TGC ATG GTT TCC TTT TTC AAT CTG GAT GGG ATG GAG CAC	192
Asp Gly Ala Cys Met Val Ser Phe Phe Asn Leu Asp Gly Met Glu His	
50 55 60	
CAT GTG CGC ACC TGC ATC CCC AAA GTG GAG CTG GTC CCT GCC GGG AAG	240
His Val Arg Thr Cys Ile Pro Lys Val Glu Leu Val Pro Ala Gly Lys	
65 70 75 80	
CCC TTC TAC TGC CTG AGC TCG GAG GAC CTG CGC AAC ACC CAC TGC TGC	288
Pro Phe Tyr Cys Leu Ser Ser Glu Asp Leu Arg Asn Thr His Cys Cys	
85 90 95	
TAC ACT GAC TAC TGC AAC AGG ATC GAC TTG AGG GTG CCC AGT GGT CAC	336
Tyr Thr Asp Tyr Cys Asn Arg Ile Asp Leu Arg Val Pro Ser Gly His	
100 105 110	
CTC AAG GAG CCT GAG CAC CCG TCC ATG TGG GGC CCG GTG GAG CTG GTA	384
Leu Lys Glu Pro Glu His Pro Ser Met Trp Gly Pro Val Glu Leu Val	
115 120 125	
GGC ATC ATC GCC GGC CCG GTG TTC CTC CTG TTC CTC ATC ATC ATC ATT	432
Gly Ile Ile Ala Gly Pro Val Phe Leu Leu Phe Leu Ile Ile Ile Ile	
130 135 140	
GTT TTC CTT GTC ATT AAC TAT CAT CAG CGT GTC TAT CAC AAC CGC CAG	480
Val Phe Leu Val Ile Asn Tyr His Gln Arg Val Tyr His Asn Arg Gln	
145 150 155 160	
AGA CTG GAC ATG GAA GAT CCC TCA TGT GAG ATG TGT CTC TCC AAA GAC	528
Arg Leu Asp Met Glu Asp Pro Ser Cys Glu Met Cys Leu Ser Lys Asp	
165 170 175	
AAG ACG CTC CAG GAT CTT GTC TAC GAT CTC TCC ACC TCA GGG TCT GGC	576
Lys Thr Leu Gln Asp Leu Val Tyr Asp Leu Ser Thr Ser Gly Ser Gly	
180 185 190	
TCA GGG TTA CCC CTC TTT GTC CAG CGC ACA GTG GCC CGA ACC ATC GTT	624
Ser Gly Leu Pro Leu Phe Val Gln Arg Thr Val Ala Arg Thr Ile Val	
195 200 205	
TTA CAA GAG ATT ATT GGC AAG GGT CGG TTT GGG GAA GTA TGG CGG GGC	672
Leu Gln Glu Ile Ile Gly Lys Gly Arg Phe Gly Glu Val Trp Arg Gly	
210 215 220	

CGC TGG AGG GGT GGT GAT GTG GCT GTG AAA ATA TTC TCT TCT CGT GAA Arg Trp Arg Gly Gly Asp Val Ala Val Lys Ile Phe Ser Ser Arg Glu 225 230 235 240	720
GAA CGG TCT TGG TTC AGG GAA GCA GAG ATA TAC CAG ACG GTC ATG CTG Glu Arg Ser Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu 245 250 255	768
CGC CAT GAA AAC ATC CTT GGA TTT ATT GCT GCT GAC AAT AAA GAT AAT Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn 260 265 270	816
GGC ACC TGG ACA CAG CTG TGG CTT GTT TCT GAC TAT CAT GAG CAC GGG Gly Thr Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly 275 280 285	864
TCC CTG TTT GAT TAT CTG AAC CGG TAC ACA GTG ACA ATT GAG GGG ATG Ser Leu Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Ile Glu Gly Met 290 295 300	912
ATT AAG CTG GCC TTG TCT GCT GCT AGT GGG CTG GCA CAC CTG CAC ATG Ile Lys Leu Ala Leu Ser Ala Ala Ser Gly Leu Ala His Leu His Met 305 310 315 320	960
GAG ATC GTG GGC ACC CAA GGG AAG CCT GGA ATT GCT CAT CGA GAC TTA Glu Ile Val Gly Thr Gln Gly Lys Pro Gly Ile Ala His Arg Asp Leu 325 330 335	1008
AAG TCA AAG AAC ATT CTG GTG AAG AAA AAT GGC ATG TGT GCC ATA GCA Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Met Cys Ala Ile Ala 340 345 350	1056
GAC CTG GGC CTG GCT GTC CGT CAT GAT GCA GTC ACT GAC ACC ATT GAC Asp Leu Gly Leu Ala Val Arg His Asp Ala Val Thr Asp Thr Ile Asp 355 360 365	1104
ATT GCC CCG AAT CAG AGG GTG GGG ACC AAA CGA TAC ATG GCC CCT GAA Ile Ala Pro Asn Gln Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu 370 375 380	1152
GTA CTT GAT GAA ACC ATT AAT ATG AAA CAC TTT GAC TCC TTT AAA TGT Val Leu Asp Glu Thr Ile Asn Met Lys His Phe Asp Ser Phe Lys Cys 385 390 395 400	1200
GCT GAT ATT TAT GCC CTC GGG CTT GTA TAT TGG GAG ATT GCT CGA AGA Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp Glu Ile Ala Arg Arg 405 410 415	1248
TGC AAT TCT GGA GGA GTC CAT GAA GAA TAT CAG CTG CCA TAT TAC GAC Cys Asn Ser Gly Gly Val His Glu Glu Tyr Gln Leu Pro Tyr Tyr Asp 420 425 430	1296
TTA GTG CCC TCT GAC CCT TCC ATT GAG GAA ATG CGA AAG GTT GTA TGT Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met Arg Lys Val Val Cys 435 440 445	1344
GAT CAG AAG CTG CGT CCC AAC ATC CCC AAC TGG TGG CAG AGT TAT GAG Asp Gln Lys Leu Arg Pro Asn Ile Pro Asn Trp Trp Gln Ser Tyr Glu 450 455 460	1392

GCA CTG CGG GTG ATG GGG AAG ATG ATG CGA GAG TGT TGG TAT GCC AAC	1440
Ala Leu Arg Val Met Gly Lys Met Met Arg Glu Cys Trp Tyr Ala Asn	
465 470 475 480	
GGC GCA GCC CGC CTG ACG GCC CTG CGC ATC AAG AAG ACC CTC TCC CAG	1488
Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ser Gln	
485 490 495	
CTC AGC GTG CAG GAA GAC GTG AAG ATC TAACTGCTCC CTCTCTCCAC	1535
Leu Ser Val Gln Glu Asp Val Lys Ile	
500 505	
ACGGAGCTCC TGGCAGCGAG AACTACGCAC AGCTGCCGCG TTGAGCGTAC GATGGAGGCC	1595
TACCTCTCGT TTCTGCCCAG CCCTCTGTGG CCAGGAGCCC TGGCCCGCAA GAGGGACAGA	1655
GCCCCGGAGA GACTCGCTCA CTCCCATGTT GGGTTTGAGA CAGACACCTT TTCTATTTAC	1715
CTCCTAATGG CATGGAGACT CTGAGAGCGA ATTGTGTGGA GAACTCAGTG CCACACCTCG	1775
AACTGGTTGT AGTGGGAAGT CCGCGGAAAC CCGGTGCATC TGGCACGTGG CCAGGAGCCA	1835
TGACAGGGGC GCTTGGGAGG GGCCGGAGGA ACCGAGGTGT TGCCAGTGCT AAGCTGCCCT	1895
GAGGGTTTCC TTCGGGGACC AGCCCACAGC ACACCAAGGT GGCCCGGAAG AACCAGAAGT	1955
GCAGCCCCTC TCACAGGCAG CTCTGAGCCG CGCTTTCCCC TCCTCCCTGG GATGGACGCT	2015
GCCGGGAGAC TGCCAGTGGA GACGGAATCT GCCGCTTTGT CTGTCCAGCC GTGTGTGCAT	2075
GTGCCGAGGT GCGTCCCCCG TTGTGCCTGG TTCGTGCCAT GCCCTTACAC GTGCGTGTGA	2135
GTGTGTGTGT GTGTCTGTAG GTGCGCACTT ACCTGCTTGA GCTTTCTGTG CATGTGCAGG	2195
TCGGGGGTGT GGTCGTCATG CTGTCCGTGC TTGCTGGTGC CTCTTTTCAG TAGTGAGCAG	2255
CATCTAGTTT CCCTGGTGCC CTTCCCTGGA GGTCTCTCCC TCCCCAGAG CCCCTCATGC	2315
CACAGTGGTA CTCTGTGT	2333

## (2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 505 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met	Ala	Glu	Ser	Ala	Gly	Ala	Ser	Ser	Phe	Phe	Pro	Leu	Val	Val	Leu
1				5					10					15	
Leu	Leu	Ala	Gly	Ser	Gly	Gly	Ser	Gly	Pro	Arg	Gly	Val	Gln	Ala	Leu
			20					25						30	

54

Leu Cys Ala Cys Thr Ser Cys Leu Gln Ala Asn Tyr Thr Cys Glu Thr  
 35 40 45  
 Asp Gly Ala Cys Met Val Ser Phe Phe Asn Leu Asp Gly Met Glu His  
 50 55 60  
 His Val Arg Thr Cys Ile Pro Lys Val Glu Leu Val Pro Ala Gly Lys  
 65 70 75 80  
 Pro Phe Tyr Cys Leu Ser Ser Glu Asp Leu Arg Asn Thr His Cys Cys  
 85 90 95  
 Tyr Thr Asp Tyr Cys Asn Arg Ile Asp Leu Arg Val Pro Ser Gly His  
 100 105 110  
 Leu Lys Glu Pro Glu His Pro Ser Met Trp Gly Pro Val Glu Leu Val  
 115 120 125  
 Gly Ile Ile Ala Gly Pro Val Phe Leu Leu Phe Leu Ile Ile Ile Ile  
 130 135 140  
 Val Phe Leu Val Ile Asn Tyr His Gln Arg Val Tyr His Asn Arg Gln  
 145 150 155 160  
 Arg Leu Asp Met Glu Asp Pro Ser Cys Glu Met Cys Leu Ser Lys Asp  
 165 170 175  
 Lys Thr Leu Gln Asp Leu Val Tyr Asp Leu Ser Thr Ser Gly Ser Gly  
 180 185 190  
 Ser Gly Leu Pro Leu Phe Val Gln Arg Thr Val Ala Arg Thr Ile Val  
 195 200 205  
 Leu Gln Glu Ile Ile Gly Lys Gly Arg Phe Gly Glu Val Trp Arg Gly  
 210 215 220  
 Arg Trp Arg Gly Gly Asp Val Ala Val Lys Ile Phe Ser Ser Arg Glu  
 225 230 235 240  
 Glu Arg Ser Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu  
 245 250 255  
 Arg His Glu Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn  
 260 265 270  
 Gly Thr Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly  
 275 280 285  
 Ser Leu Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Ile Glu Gly Met  
 290 295 300  
 Ile Lys Leu Ala Leu Ser Ala Ala Ser Gly Leu Ala His Leu His Met  
 305 310 315 320  
 Glu Ile Val Gly Thr Gln Gly Lys Pro Gly Ile Ala His Arg Asp Leu  
 325 330 335  
 Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Met Cys Ala Ile Ala  
 340 345 350



55

Asp Leu Gly Leu Ala Val Arg His Asp Ala Val Thr Asp Thr Ile Asp  
 355 360 365  
 Ile Ala Pro Asn Gln Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu  
 370 375 380  
 Val Leu Asp Glu Thr Ile Asn Met Lys His Phe Asp Ser Phe Lys Cys  
 385 390 395 400  
 Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp Glu Ile Ala Arg Arg  
 405 410 415  
 Cys Asn Ser Gly Gly Val His Glu Glu Tyr Gln Leu Pro Tyr Tyr Asp  
 420 425 430  
 Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met Arg Lys Val Val Cys  
 435 440 445  
 Asp Gln Lys Leu Arg Pro Asn Ile Pro Asn Trp Trp Gln Ser Tyr Glu  
 450 455 460  
 Ala Leu Arg Val Met Gly Lys Met Met Arg Glu Cys Trp Tyr Ala Asn  
 465 470 475 480  
 Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ser Gln  
 485 490 495  
 Leu Ser Val Gln Glu Asp Val Lys Ile  
 500 505

## (2) INFORMATION FOR SEQ ID NO: 9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2308 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 77..1585

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

GGCGAGGCCGA GGTTCGCTGG GGTGAGGCAG CGGCGCGGCC GGGCCGGGCC GGGCCACAGG

60

56

CGGTGGCGGC GGGACC ATG GAG GCG GCG GTC GCT GCT CCG CGT CCC CGG	109
Met Glu Ala Ala Val Ala Ala Pro Arg Pro Arg	
1 5 10	
CTG CTC CTC CTC GTG CTG GCG GCG GCG GCG GCG GCG GCG GCG CTG	157
Leu Leu Leu Leu Val Leu Ala Ala Ala Ala Ala Ala Ala Ala Leu	
15 20 25	
CTC CCG GGG GCG ACG GCG TTA CAG TGT TTC TGC CAC CTC TGT ACA AAA	205
Leu Pro Gly Ala Thr Ala Leu Gln Cys Phe Cys His Leu Cys Thr Lys	
30 35 40	
GAC AAT TTT ACT TGT GTG ACA GAT GGG CTC TGC TTT GTC TCT GTC ACA	253
Asp Asn Phe Thr Cys Val Thr Asp Gly Leu Cys Phe Val Ser Val Thr	
45 50 55	
GAG ACC ACA GAC AAA GTT ATA CAC AAC AGC ATG TGT ATA GCT GAA ATT	301
Glu Thr Thr Asp Lys Val Ile His Asn Ser Met Cys Ile Ala Glu Ile	
60 65 70 75	
GAC TTA ATT CCT CGA GAT AGG CCG TTT GTA TGT GCA CCC TCT TCA AAA	349
Asp Leu Ile Pro Arg Asp Arg Pro Phe Val Cys Ala Pro Ser Ser Lys	
80 85 90	
ACT GGG TCT GTG ACT ACA ACA TAT TGC TGC AAT CAG GAC CAT TGC AAT	397
Thr Gly Ser Val Thr Thr Thr Tyr Cys Cys Asn Gln Asp His Cys Asn	
95 100 105	
AAA ATA GAA CTT CCA ACT ACT GTA AAG TCA TCA CCT GGC CTT GGT CCT	445
Lys Ile Glu Leu Pro Thr Thr Val Lys Ser Ser Pro Gly Leu Gly Pro	
110 115 120	
GTG GAA CTG GCA GCT GTC ATT GCT GGA CCA GTG TGC TTC GTC TGC ATC	493
Val Glu Leu Ala Ala Val Ile Ala Gly Pro Val Cys Phe Val Cys Ile	
125 130 135	
TCA CTC ATG TTG ATG GTC TAT ATC TGC CAC AAC CGC ACT GTC ATT CAC	541
Ser Leu Met Leu Met Val Tyr Ile Cys His Asn Arg Thr Val Ile His	
140 145 150 155	
CAT CGA GTG CCA AAT GAA GAG GAC CCT TCA TTA GAT CGC CCT TTT ATT	589
His Arg Val Pro Asn Glu Glu Asp Pro Ser Leu Asp Arg Pro Phe Ile	
160 165 170	
TCA GAG GGT ACT ACG TTG AAA GAC TTA ATT TAT GAT ATG ACA ACG TCA	637
Ser Glu Gly Thr Thr Leu Lys Asp Leu Ile Tyr Asp Met Thr Thr Ser	
175 180 185	
GGT TCT GGC TCA GGT TTA CCA TTG CTT GTT CAG AGA ACA ATT GCG AGA	685
Gly Ser Gly Ser Gly Leu Pro Leu Leu Val Gln Arg Thr Ile Ala Arg	
190 195 200	
ACT ATT GTG TTA CAA GAA AGC ATT GGC AAA GGT CGA TTT GGA GAA GTT	733
Thr Ile Val Leu Gln Glu Ser Ile Gly Lys Gly Arg Phe Gly Glu Val	
205 210 215	
TGG AGA GGA AAG TGG CCG GGA GAA GAA GTT GCT GTT AAG ATA TTC TCC	781
Trp Arg Gly Lys Trp Arg Gly Glu Glu Val Ala Val Lys Ile Phe Ser	
220 225 230 235	

TCT	AGA	GAA	GAA	CGT	TCG	TGG	TTC	CGT	GAG	GCA	GAG	ATT	TAT	CAA	ACT	829
Ser	Arg	Glu	Glu	Arg	Ser	Trp	Phe	Arg	Glu	Ala	Glu	Ile	Tyr	Gln	Thr	
				240					245					250		
GTA	ATG	TTA	CGT	CAT	GAA	AAC	ATC	CTG	GGA	TTT	ATA	GCA	GCA	GAC	AAT	877
Val	Met	Leu	Arg	His	Glu	Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	Asp	Asn	
			255					260					265			
AAA	GAC	AAT	GGT	ACT	TGG	ACT	CAG	CTC	TGG	TTG	GTG	TCA	GAT	TAT	CAT	925
Lys	Asp	Asn	Gly	Thr	Trp	Thr	Gln	Leu	Trp	Leu	Val	Ser	Asp	Tyr	His	
		270					275					280				
GAG	CAT	GGA	TCC	CTT	TTT	GAT	TAC	TTA	AAC	AGA	TAC	ACA	GTT	ACT	GTG	973
Glu	His	Gly	Ser	Leu	Phe	Asp	Tyr	Leu	Asn	Arg	Tyr	Thr	Val	Thr	Val	
	285					290					295					
GAA	GGA	ATG	ATA	AAA	CTT	GCT	CTG	TCC	ACG	GCG	AGC	GGT	CTT	GCC	CAT	1021
Glu	Gly	Met	Ile	Lys	Leu	Ala	Leu	Ser	Thr	Ala	Ser	Gly	Leu	Ala	His	
300				305					310					315		
CTT	CAC	ATG	GAG	ATT	GTT	GGT	ACC	CAA	GGA	AAG	CCA	GCC	ATT	GCT	CAT	1069
Leu	His	Met	Glu	Ile	Val	Gly	Thr	Gln	Gly	Lys	Pro	Ala	Ile	Ala	His	
			320					325					330			
AGA	GAT	TTG	AAA	TCA	AAG	AAT	ATC	TTG	GTA	AAG	AAG	AAT	GGA	ACT	TGC	1117
Arg	Asp	Leu	Lys	Ser	Lys	Asn	Ile	Leu	Val	Lys	Lys	Asn	Gly	Thr	Cys	
			335					340					345			
TGT	ATT	GCA	GAC	TTA	GGA	CTG	GCA	GTA	AGA	CAT	GAT	TCA	GCC	ACA	GAT	1165
Cys	Ile	Ala	Asp	Leu	Gly	Leu	Ala	Val	Arg	His	Asp	Ser	Ala	Thr	Asp	
	350					355					360					
ACC	ATT	GAT	ATT	GCT	CCA	AAC	CAC	AGA	GTG	GGA	ACA	AAA	AGG	TAC	ATG	1213
Thr	Ile	Asp	Ile	Ala	Pro	Asn	His	Arg	Val	Gly	Thr	Lys	Arg	Tyr	Met	
	365			370							375					
GCC	CCT	GAA	GTT	CTC	GAT	GAT	TCC	ATA	AAT	ATG	AAA	CAT	TTT	GAA	TCC	1261
Ala	Pro	Glu	Val	Leu	Asp	Asp	Ser	Ile	Asn	Met	Lys	His	Phe	Glu	Ser	
380				385					390					395		
TTC	AAA	CGT	GCT	GAC	ATC	TAT	GCA	ATG	GGC	TTA	GTA	TTC	TGG	GAA	ATT	1309
Phe	Lys	Arg	Ala	Asp	Ile	Tyr	Ala	Met	Gly	Leu	Val	Phe	Trp	Glu	Ile	
			400					405					410			
GCT	CGA	CGA	TGT	TCC	ATT	GGT	GGA	ATT	CAT	GAA	GAT	TAC	CAA	CTG	CCT	1357
Ala	Arg	Arg	Cys	Ser	Ile	Gly	Gly	Ile	His	Glu	Asp	Tyr	Gln	Leu	Pro	
			415				420						425			
TAT	TAT	GAT	CTT	GTA	CCT	TCT	GAC	CCA	TCA	GTT	GAA	GAA	ATG	AGA	AAA	1405
Tyr	Tyr	Asp	Leu	Val	Pro	Ser	Asp	Pro	Ser	Val	Glu	Glu	Met	Arg	Lys	
		430				435					440					
GTT	GTT	TGT	GAA	CAG	AAG	TTA	AGG	CCA	AAT	ATC	CCA	AAC	AGA	TGG	CAG	1453
Val	Val	Cys	Glu	Gln	Lys	Leu	Arg	Pro	Asn	Ile	Pro	Asn	Arg	Trp	Gln	
	445					450					455					
AGC	TGT	GAA	GCC	TTG	AGA	GTA	ATG	GCT	AAA	ATT	ATG	AGA	GAA	TGT	TGG	1501
Ser	Cys	Glu	Ala	Leu	Arg	Val	Met	Ala	Lys	Ile	Met	Arg	Glu	Cys	Trp	
460				465					470					475		

58

TAT GCC AAT CGA GCA GCT AGG CTT ACA GCA TTG CGG ATT AAG AAA ACA	1549
Tyr Ala Asn Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr	
480 485 490	
TTA TCG CAA CTC AGT CAA CAG GAA GGC ATC AAA ATG TAATTCTACA	1595
Leu Ser Gln Leu Ser Gln Gln Glu Gly Ile Lys Met	
495 500	
GCTTGCCTG AACTCTCCTT TTTCTTCAG ATCTGCTCCT GGGTTTAAAT TTGGGAGGTC	1655
AGTTGTTCTA CCTCACTGAG AGGGAACAGA AGGATATTGC TTCCTTTTGC AGCAGTGTA	1715
TAAAGTCAAT TAAAACTTC CCAGGATTTT TTTGGACCCA GGAAACAGCC ATCTGGGTCC	1775
TTTCTGTGCA CTATGAACGC TTCTTTCCCA GGACAGAAAA TGTGTAGTCT ACCTTTATTT	1835
TTTATTAACA AACTTGTTT TTTAAAAAGA TGATTGCTGG TCTTAACTTT AGGTAACCTCT	1895
GCTGTGCTGG AGATCATCTT TAAGGGCAAA GGAGTTGGAT TGCTGAATTA CAATGAAACA	1955
TGTCTTATTA CTAAAGAAAG TGATTTACTC CTGGTTAGTA CATTCTCAGA GGATTCTGAA	2015
CCACTAGAGT TTCCTTGATT CAGACTTTGA ATGTACTGTT CTATAGTTTT TCAGGATCTT	2075
AAACTAACA CTTATAAAAC TCTTATCTTG AGTCTAAAAA TGACCTCATA TAGTAGTGAG	2135
GAACATAATT CATGCAATTG TATTTTGTAT ACTATTATTG TTCTTTCCTT TATTCAGAAC	2195
ATTACATGCC TTCAAATGG GATTGTACTA TACCAGTAAG TGCCACTTCT GTGTCTTTCT	2255
AATGGAAATG AGTAGAATTG CTGAAAGTCT CTATGTTAAA ACCTATAGTG TTT	2308

## (2) INFORMATION FOR SEQ ID NO: 10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 503 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met	Glu	Ala	Ala	Val	Ala	Ala	Pro	Arg	Pro	Arg	Leu	Leu	Leu	Leu	Val
1				5					10					15	
Leu	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Ala	Leu	Leu	Pro	Gly	Ala
			20				25							30	Thr
Ala	Leu	Gln	Cys	Phe	Cys	His	Leu	Cys	Thr	Lys	Asp	Asn	Phe	Thr	Cys
			35				40						45		
Val	Thr	Asp	Gly	Leu	Cys	Phe	Val	Ser	Val	Thr	Glu	Thr	Thr	Asp	Lys
			50			55					60				
Val	Ile	His	Asn	Ser	Met	Cys	Ile	Ala	Glu	Ile	Asp	Leu	Ile	Pro	Arg
65					70					75				80	

Asp Arg Pro Phe Val Cys Ala Pro Ser Ser Lys Thr Gly Ser Val Thr  
 85 90 95  
 Thr Thr Tyr Cys Cys Asn Gln Asp His Cys Asn Lys Ile Glu Leu Pro  
 100 105 110  
 Thr Thr Val Lys Ser Ser Pro Gly Leu Gly Pro Val Glu Leu Ala Ala  
 115 120 125  
 Val Ile Ala Gly Pro Val Cys Phe Val Cys Ile Ser Leu Met Leu Met  
 130 135 140  
 Val Tyr Ile Cys His Asn Arg Thr Val Ile His His Arg Val Pro Asn  
 145 150 155 160  
 Glu Glu Asp Pro Ser Leu Asp Arg Pro Phe Ile Ser Glu Gly Thr Thr  
 165 170 175  
 Leu Lys Asp Leu Ile Tyr Asp Met Thr Thr Ser Gly Ser Gly Ser Gly  
 180 185 190  
 Leu Pro Leu Leu Val Gln Arg Thr Ile Ala Arg Thr Ile Val Leu Gln  
 195 200 205  
 Glu Ser Ile Gly Lys Gly Arg Phe Gly Glu Val Trp Arg Gly Lys Trp  
 210 215 220  
 Arg Gly Glu Glu Val Ala Val Lys Ile Phe Ser Ser Arg Glu Glu Arg  
 225 230 235 240  
 Ser Trp Phe Arg Glu Ala Glu Ile Tyr Gln Thr Val Met Leu Arg His  
 245 250 255  
 Glu Asn Ile Leu Gly Phe Ile Ala Ala Asp Asn Lys Asp Asn Gly Thr  
 260 265 270  
 Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly Ser Leu  
 275 280 285  
 Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Val Glu Gly Met Ile Lys  
 290 295 300  
 Leu Ala Leu Ser Thr Ala Ser Gly Leu Ala His Leu His Met Glu Ile  
 305 310 315 320  
 Val Gly Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser  
 325 330 335  
 Lys Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu  
 340 345 350  
 Gly Leu Ala Val Arg His Asp Ser Ala Thr Asp Thr Ile Asp Ile Ala  
 355 360 365  
 Pro Asn His Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu  
 370 375 380  
 Asp Asp Ser Ile Asn Met Lys His Phe Glu Ser Phe Lys Arg Ala Asp  
 385 390 395 400

60

Ile	Tyr	Ala	Met	Gly	Leu	Val	Phe	Trp	Glu	Ile	Ala	Arg	Arg	Cys	Ser			
									410								415	
Ile	Gly	Gly	Ile	His	Glu	Asp	Tyr	Gln	Leu	Pro	Tyr	Tyr	Asp	Leu	Val			
								425								430		
Pro	Ser	Asp	Pro	Ser	Val	Glu	Glu	Met	Arg	Lys	Val	Val	Cys	Glu	Gln			
						435						440					445	
Lys	Leu	Arg	Pro	Asn	Ile	Pro	Asn	Arg	Trp	Gln	Ser	Cys	Glu	Ala	Leu			
					450						455						460	
Arg	Val	Met	Ala	Lys	Ile	Met	Arg	Glu	Cys	Trp	Tyr	Ala	Asn	Gly	Ala			
				465						470					475			480
Ala	Arg	Leu	Thr	Ala	Leu	Arg	Ile	Lys	Lys	Thr	Leu	Ser	Gln	Leu	Ser			
					485						490						495	
Gln	Gln	Glu	Gly	Ile	Lys	Met												
				500														

(2) INFORMATION FOR SEQ ID NO: 11:

**(i) SEQUENCE CHARACTERISTICS:**

- (A) LENGTH: 1922 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: unknown  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) **FEATURE:**

(A) NAME/KEY: CDS

(B) LOCATION: 241..1746

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GAGAGCACAG	CCCTTCCCAG	TCCCCGGAGC	CGCGCGGCCA	CGCGCGCATG	ATCAAGACCT	60
TTTCCCCGGC	CCCACAGGGC	CTCTGGACGT	GAGACCCCGG	CCGCCTCCGC	AAGGAGAGGC	120
GGGGGTCGAG	TCGCCCTGTC	CAAAGGCCTC	AATCTAAACA	ATCTTGATTC	CTGTTGCCGG	180
CTGGCGGGAC	CCTGAATGGC	AGGAAATCTC	ACCACATCTC	TTCTCCTATC	TCCAAGGACC	240
ATG ACC TTG GGG AGC TTC AGA AGG GGC CTT TTG ATG CTG TCG GTG GCC	288					
Met Thr Leu Gly Ser Phe Arg Arg Gly Leu Leu Met Leu Ser Val Ala						
1 5 10 15						

TTG GGC CTA ACC CAG GGG AGA CTT GCG AAG CCT TCC AAG CTG GTG AAC	336
Leu Gly Leu Thr Gln Gly Arg Leu Ala Lys Pro Ser Lys Leu Val Asn	
20 25 30	
TGC ACT TGT GAG AGC CCA CAC TGC AAG AGA CCA TTC TGC CAG GGG TCA	384
Cys Thr Cys Glu Ser Pro His Cys Lys Arg Pro Phe Cys Gln Gly Ser	
35 40 45	
TGG TGC ACA GTG GTG CTG GTT CGA GAG CAG GGC AGG CAC CCC CAG GTC	432
Trp Cys Thr Val Val Leu Val Arg Glu Gln Gly Arg His Pro Gln Val	
50 55 60	
TAT CCG GGC TGT GGG AGC CTG AAC CAG GAG CTC TGC TTG GGA CGT CCC	480
Tyr Arg Gly Cys Gly Ser Leu Asn Gln Glu Leu Cys Leu Gly Arg Pro	
65 70 75 80	
ACG GAG TTT CTG AAC CAT CAC TGC TGC TAT AGA TCC TTC TGC AAC CAC	528
Thr Glu Phe Leu Asn His His Cys Cys Tyr Arg Ser Phe Cys Asn His	
85 90 95	
AAC GTG TCT CTG ATG CTG GAG GCC ACC CAA ACT CCT TCG GAG GAG CCA	576
Asn Val Ser Leu Met Leu Glu Ala Thr Gln Thr Pro Ser Glu Glu Pro	
100 105 110	
GAA GTT GAT GCC CAT CTG CCT CTG ATC CTG GGT CCT GTG CTG GCC TTG	624
Glu Val Asp Ala His Leu Pro Leu Ile Leu Gly Pro Val Leu Ala Leu	
115 120 125	
CCG GTC CTG GTG GCC CTG GGT GCT CTG GGC TTG TGG CGT GTC CCG CCG	672
Pro Val Leu Val Ala Leu Gly Ala Leu Gly Leu Trp Arg Val Arg Arg	
130 135 140	
AGG CAG GAG AAG CAG CCG GAT TTG CAC AGT GAC CTG GGC GAG TCC AGT	720
Arg Gln Glu Lys Gln Arg Asp Leu His Ser Asp Leu Gly Glu Ser Ser	
145 150 155 160	
CTC ATC CTG AAG GCA TCT GAA CAG GCA GAC AGC ATG TTG GGG GAC TTC	768
Leu Ile Leu Lys Ala Ser Glu Gln Ala Asp Ser Met Leu Gly Asp Phe	
165 170 175	
CTG GAC AGC GAC TGT ACC ACG GGC AGC GGC TCG GGG CTC CCC TTC TTG	816
Leu Asp Ser Asp Cys Thr Thr Gly Ser Gly Ser Gly Leu Pro Phe Leu	
180 185 190	
GTG CAG AGG ACG GTA GCT CCG CAG GTT GCG CTG GTA GAG TGT GTG GGA	864
Val Gln Arg Thr Val Ala Arg Gln Val Ala Leu Val Glu Cys Val Gly	
195 200 205	
AAG GGC CGA TAT GGC GAG GTG TGG CGC GGT TCG TGG CAT GGC GAA AGC	912
Lys Gly Arg Tyr Gly Glu Val Trp Arg Gly Ser Trp His Gly Glu Ser	
210 215 220	
GTG GCG GTC AAG ATT TTC TCC TCA CGA GAT GAG CAG TCC TGG TTC CCG	960
Val Ala Val Lys Ile Phe Ser Ser Arg Asp Glu Gln Ser Trp Phe Arg	
225 230 235 240	
GAG ACG GAG ATC TAC AAC ACA GTT CTG CTT AGA CAC GAC AAC ATC CTA	1008
Glu Thr Glu Ile Tyr Asn Thr Val Leu Leu Arg His Asp Asn Ile Leu	
245 250 255	

GGC	TTC	ATC	GCC	TCC	GAC	ATG	ACT	TCG	CGG	AAC	TCG	AGC	ACG	CAG	CTG	1056
Gly	Phe	Ile	Ala	Ser	Asp	Met	Thr	Ser	Arg	Asn	Ser	Ser	Thr	Gln	Leu	
			260					265					270			
TGG	CTC	ATC	ACC	CAC	TAC	CAT	GAA	CAC	GGC	TCC	CTC	TAT	GAC	TTT	CTG	1104
Trp	Leu	Ile	Thr	His	Tyr	His	Glu	His	Gly	Ser	Leu	Tyr	Asp	Phe	Leu	
		275					280					285				
CAG	AGG	CAG	ACG	CTG	GAG	CCC	CAG	TTG	GCC	CTG	AGG	CTA	GCT	GTG	TCC	1152
Gln	Arg	Gln	Thr	Leu	Glu	Pro	Gln	Leu	Ala	Leu	Arg	Leu	Ala	Val	Ser	
	290					295					300					
CCG	GCC	TGC	GGC	CTG	GCG	CAC	CTA	CAT	GTG	GAG	ATC	TTT	GGC	ACT	CAA	1200
Pro	Ala	Cys	Gly	Leu	Ala	His	Leu	His	Val	Glu	Ile	Phe	Gly	Thr	Gln	
305				310						315					320	
GGC	AAA	CCA	GCC	ATT	GCC	CAT	CGT	GAC	CTC	AAG	AGT	CGC	AAT	GTG	CTG	1248
Gly	Lys	Pro	Ala	Ile	Ala	His	Arg	Asp	Leu	Lys	Ser	Arg	Asn	Val	Leu	
			325					330						335		
GTC	AAG	AGT	AAC	TTG	CAG	TGT	TGC	ATT	GCA	GAC	CTG	GGA	CTG	GCT	GTG	1296
Val	Lys	Ser	Asn	Leu	Gln	Cys	Cys	Ile	Ala	Asp	Leu	Gly	Leu	Ala	Val	
			340					345					350			
ATG	CAC	TCA	CAA	AGC	AAC	GAG	TAC	CTG	GAT	ATC	GGC	AAC	ACA	CCC	CGA	1344
Met	His	Ser	Gln	Ser	Asn	Glu	Tyr	Leu	Asp	Ile	Gly	Asn	Thr	Pro	Arg	
		355					360					365				
GTG	GGT	ACC	AAA	AGA	TAC	ATG	GCA	CCC	GAG	GTG	CTG	GAT	GAG	CAC	ATC	1392
Val	Gly	Thr	Lys	Arg	Tyr	Met	Ala	Pro	Glu	Val	Leu	Asp	Glu	His	Ile	
	370					375					380					
CGC	ACA	GAC	TGC	TTT	GAG	TCG	TAC	AAG	TGG	ACA	GAC	ATC	TGG	GCC	TTT	1440
Arg	Thr	Asp	Cys	Phe	Glu	Ser	Tyr	Lys	Trp	Thr	Asp	Ile	Trp	Ala	Phe	
385				390					395					400		
GGC	CTA	GTG	CTA	TGG	GAG	ATC	GCC	CGG	CGG	ACC	ATC	ATC	AAT	GGC	ATT	1488
Gly	Leu	Val	Leu	Trp	Glu	Ile	Ala	Arg	Arg	Thr	Ile	Ile	Asn	Gly	Ile	
			405					410						415		
GTG	GAG	GAT	TAC	AGG	CCA	CCT	TTC	TAT	GAC	ATG	GTA	CCC	AAT	GAC	CCC	1536
Val	Glu	Asp	Tyr	Arg	Pro	Pro	Phe	Tyr	Asp	Met	Val	Pro	Asn	Asp	Pro	
			420					425					430			
AGT	TTT	GAG	GAC	ATG	AAA	AAG	GTG	GTG	TGC	GTT	GAC	CAG	CAG	ACA	CCC	1584
Ser	Phe	Glu	Asp	Met	Lys	Lys	Val	Val	Cys	Val	Asp	Gln	Gln	Thr	Pro	
		435					440					445				
ACC	ATC	CCT	AAC	CGG	CTG	GCT	GCA	GAT	CCG	GTC	CTC	TCC	GGG	CTG	GCC	1632
Thr	Ile	Pro	Asn	Arg	Leu	Ala	Ala	Asp	Pro	Val	Leu	Ser	Gly	Leu	Ala	
	450					455					460					
CAG	ATG	ATG	AGA	GAG	TGC	TGG	TAC	CCC	AAC	CCC	TCT	GCT	CGC	CTC	ACC	1680
Gln	Met	Met	Arg	Glu	Cys	Trp	Tyr	Pro	Asn	Pro	Ser	Ala	Arg	Leu	Thr	
465				470						475				480		
GCA	CTG	CGC	ATA	AAG	AAG	ACA	TTG	CAG	AAG	CTC	AGT	CAC	AAT	CCA	GAG	1728
Ala	Leu	Arg	Ile	Lys	Lys	Thr	Leu	Gln	Lys	Leu	Ser	His	Asn	Pro	Glu	
			485					490						495		



AAG CCC AAA GTG ATT CAC TAGCCCAGGG CCACCAGGCT TCCTCTGCCT 1776  
 Lys Pro Lys Val Ile His  
 500

AAAGTGTGTG CTGGGGAAGA AGACATAGCC TGTCTGGGTA GAGGGAGTGA AGAGAGTGTG 1836  
 CACGCTGCCC TGTGTGTGCC TGCTCAGCTT GCTCCCAGCC CATCCAGCCA AAAATACAGC 1896  
 TGAGCTGAAA TTCAAAAAA AAAAAA 1922

## (2) INFORMATION FOR SEQ ID NO: 12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 502 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Met	Thr	Leu	Gly	Ser	Phe	Arg	Arg	Gly	Leu	Leu	Met	Leu	Ser	Val	Ala
1				5					10					15	
Leu	Gly	Leu	Thr	Gln	Gly	Arg	Leu	Ala	Lys	Pro	Ser	Lys	Leu	Val	Asn
		20						25					30		
Cys	Thr	Cys	Glu	Ser	Pro	His	Cys	Lys	Arg	Pro	Phe	Cys	Gln	Gly	Ser
		35					40					45			
Trp	Cys	Thr	Val	Val	Leu	Val	Arg	Glu	Gln	Gly	Arg	His	Pro	Gln	Val
	50					55					60				
Tyr	Arg	Gly	Cys	Gly	Ser	Leu	Asn	Gln	Glu	Leu	Cys	Leu	Gly	Arg	Pro
65					70					75					80
Thr	Glu	Phe	Leu	Asn	His	His	Cys	Cys	Tyr	Arg	Ser	Phe	Cys	Asn	His
				85					90					95	
Asn	Val	Ser	Leu	Met	Leu	Glu	Ala	Thr	Gln	Thr	Pro	Ser	Glu	Glu	Pro
			100					105					110		
Glu	Val	Asp	Ala	His	Leu	Pro	Leu	Ile	Leu	Gly	Pro	Val	Leu	Ala	Leu
		115					120					125			
Pro	Val	Leu	Val	Ala	Leu	Gly	Ala	Leu	Gly	Leu	Trp	Arg	Val	Arg	Arg
	130					135					140				
Arg	Gln	Glu	Lys	Gln	Arg	Asp	Leu	His	Ser	Asp	Leu	Gly	Glu	Ser	Ser
145					150					155					160
Leu	Ile	Leu	Lys	Ala	Ser	Glu	Gln	Ala	Asp	Ser	Met	Leu	Gly	Asp	Phe
			165						170					175	
Leu	Asp	Ser	Asp	Cys	Thr	Thr	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Phe	Leu
			180					185					190		

Val Gln Arg Thr Val Ala Arg Gln Val Ala Leu Val Glu Cys Val Gly  
 195 200 205  
 Lys Gly Arg Tyr Gly Glu Val Trp Arg Gly Ser Trp His Gly Glu Ser  
 210 215 220  
 Val Ala Val Lys Ile Phe Ser Ser Arg Asp Glu Gln Ser Trp Phe Arg  
 225 230 235 240  
 Glu Thr Glu Ile Tyr Acn Thr Val Leu Leu Arg His Asp Asn Ile Leu  
 245 250 255  
 Gly Phe Ile Ala Ser Asp Met Thr Ser Arg Asn Ser Ser Thr Gln Leu  
 260 265 270  
 Trp Leu Ile Thr His Tyr His Glu His Gly Ser Leu Tyr Asp Phe Leu  
 275 280 285  
 Gln Arg Gln Thr Leu Glu Pro Gln Leu Ala Leu Arg Leu Ala Val Ser  
 290 295 300  
 Pro Ala Cys Gly Leu Ala His Leu His Val Glu Ile Phe Gly Thr Gln  
 305 310 315 320  
 Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Arg Asn Val Leu  
 325 330 335  
 Val Lys Ser Asn Leu Gln Cys Cys Ile Ala Asp Leu Gly Leu Ala Val  
 340 345 350  
 Met His Ser Gln Ser Asn Glu Tyr Leu Asp Ile Gly Asn Thr Pro Arg  
 355 360 365  
 Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu His Ile  
 370 375 380  
 Arg Thr Asp Cys Phe Glu Ser Tyr Lys Trp Thr Asp Ile Trp Ala Phe  
 385 390 395 400  
 Gly Leu Val Leu Trp Glu Ile Ala Arg Arg Thr Ile Ile Asn Gly Ile  
 405 410 415  
 Val Glu Asp Tyr Arg Pro Pro Phe Tyr Asp Met Val Pro Asn Asp Pro  
 420 425 430  
 Ser Phe Glu Asp Met Lys Lys Val Val Cys Val Asp Gln Gln Thr Pro  
 435 440 445  
 Thr Ile Pro Asn Arg Leu Ala Ala Asp Pro Val Leu Ser Gly Leu Ala  
 450 455 460  
 Gln Met Met Arg Glu Cys Trp Tyr Pro Asn Pro Ser Ala Arg Leu Thr  
 465 470 475 480  
 Ala Leu Arg Ile Lys Lys Thr Leu Gln Lys Leu Ser His Asn Pro Glu  
 485 490 495  
 Lys Pro Lys Val Ile His  
 500

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2070 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 217..1812

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

ATTCATGAGA TGAAGCATA GGTCAAAGCT GTTCGGAGAA ATTGGAACTA CAGTTTTATC	60
TAGCCACATC TCTGAGAATT CTGAAGAAAG CAGCAGGTGA AAGTCATTGC CAAGTGATTT	120
TGTTCTGTAA GGAAGCCTCC CTCATTCACT TACACCAGTG AGACAGCAGG ACCAGTCATT	180
CAAAGGGCCG TGTACAGGAC GCGTGGCAAT CAGACA ATG ACT CAG CTA TAC ACT	234
Met Thr Gln Leu Tyr Thr	
1 5	
TAC ATC AGA TTA CTG GGA GCC TGT CTG TTC ATC ATT TCT CAT GTT CAA	282
Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe Ile Ile Ser His Val Gln	
10 15 20	
GGG CAG AAT CTA GAT AGT ATG CTC CAT GGC ACT GGT ATG AAA TCA GAC	330
Gly Gln Asn Leu Asp Ser Met Leu His Gly Thr Gly Met Lys Ser Asp	
25 30 35	
TTG GAC CAG AAG AAG CCA GAA AAT GGA GTG ACT TTA GCA CCA GAG GAT	378
Leu Asp Gln Lys Lys Pro Glu Asn Gly Val Thr Leu Ala Pro Glu Asp	
40 45 50	
ACC TTG CCT TTC TTA AAG TGC TAT TGC TCA GGA CAC TGC CCA GAT GAT	426
Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser Gly His Cys Pro Asp Asp	
55 60 65 70	
GCT ATT AAT AAC ACA TGC ATA ACT AAT GGC CAT TGC TTT GCC ATT ATA	474
Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly His Cys Phe Ala Ile Ile	
75 80 85	
GAA GAA GAT GAT CAG GGA GAA ACC ACA TTA ACT TCT GGG TGT ATG AAG	522
Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu Thr Ser Gly Cys Met Lys	
90 95 100	

TAT	GAA	GGC	TCT	GAT	TTT	CAA	TGC	AAG	GAT	TCA	CCG	AAA	GCC	CAG	CTA	570
Tyr	Glu	Gly	Ser	Asp	Phe	Gln	Cys	Lys	Asp	Ser	Pro	Lys	Ala	Gln	Leu	
		105					110					115				
CGC	AGG	ACA	ATA	GAA	TGT	TGT	CGG	ACC	AAT	TTG	TGC	AAC	CAG	TAT	TTG	618
Arg	Arg	Thr	Ile	Glu	Cys	Cys	Arg	Thr	Asn	Leu	Cys	Asn	Gln	Tyr	Leu	
		120				125					130					
CAG	CCT	ACA	CTG	CCC	CCT	GTT	GTT	ATA	GGT	CCG	TTC	TTT	GAT	GGC	AGC	666
Gln	Pro	Thr	Leu	Pro	Pro	Val	Val	Ile	Gly	Pro	Phe	Phe	Asp	Gly	Ser	
					140					145					150	
ATC	CGA	TGG	CTG	GTT	GTG	CTC	ATT	TCC	ATG	GCT	GTC	TGT	ATA	GTT	GCT	714
Ile	Arg	Trp	Leu	Val	Val	Leu	Ile	Ser	Met	Ala	Val	Cys	Ile	Val	Ala	
				155					160					165		
ATG	ATC	ATC	TTC	TCC	AGC	TGC	TTT	TGC	TAT	AAG	CAT	TAT	TGT	AAG	AGT	762
Met	Ile	Ile	Phe	Ser	Ser	Cys	Phe	Cys	Tyr	Lys	His	Tyr	Cys	Lys	Ser	
			170					175					180			
ATC	TCA	AGC	AGG	GGT	CGT	TAC	AAC	CGT	GAT	TTG	GAA	CAG	GAT	GAA	GCA	810
Ile	Ser	Ser	Arg	Gly	Arg	Tyr	Asn	Arg	Asp	Leu	Glu	Gln	Asp	Glu	Ala	
			185				190					195				
TTT	ATT	CCA	GTA	GGA	GAA	TCA	TTG	AAA	GAC	CTG	ATT	GAC	CAG	TCC	CAA	858
Phe	Ile	Pro	Val	Gly	Glu	Ser	Leu	Lys	Asp	Leu	Ile	Asp	Gln	Ser	Gln	
		200				205					210					
AGC	TCT	GGG	AGT	GGA	TCT	GGA	TTG	CCT	TTA	TTG	GTT	CAG	CGA	ACT	ATT	906
Ser	Ser	Gly	Ser	Gly	Ser	Gly	Leu	Pro	Leu	Leu	Val	Gln	Arg	Thr	Ile	
		215			220					225					230	
GCC	AAA	CAG	ATT	CAG	ATG	GTT	CGG	CAG	GTT	GGT	AAA	GGC	CGC	TAT	GGA	954
Ala	Lys	Gln	Ile	Gln	Met	Val	Arg	Gln	Val	Gly	Lys	Gly	Arg	Tyr	Gly	
				235					240					245		
GAA	GTA	TGG	ATG	GGT	AAA	TGG	CGT	GGT	GAA	AAA	GTG	GCT	GTC	AAA	GTG	1002
Glu	Val	Trp	Met	Gly	Lys	Trp	Arg	Gly	Glu	Lys	Val	Ala	Val	Lys	Val	
			250					255					260			
TTT	TTT	ACC	ACT	GAA	GAA	GCT	AGC	TGG	TTT	AGA	GAA	ACA	GAA	ATC	TAC	1050
Phe	Phe	Thr	Thr	Glu	Glu	Ala	Ser	Trp	Phe	Arg	Glu	Thr	Glu	Ile	Tyr	
		265					270					275				
CAG	ACG	GTG	TTA	ATG	CGT	CAT	GAA	AAT	ATA	CTT	GGT	TTT	ATA	GCT	GCA	1098
Gln	Thr	Val	Leu	Met	Arg	His	Glu	Asn	Ile	Leu	Gly	Phe	Ile	Ala	Ala	
		280				285					290					
GAC	ATT	AAA	GGC	ACT	GGT	TCC	TGG	ACT	CAG	CTG	TAT	TTG	ATT	ACT	GAT	1146
Asp	Ile	Lys	Gly	Thr	Gly	Ser	Trp	Thr	Gln	Leu	Tyr	Leu	Ile	Thr	Asp	
					300					305					310	
295																
TAC	CAT	GAA	AAT	GGA	TCT	CTC	TAT	GAC	TTC	CTG	AAA	TGT	GCC	ACA	CTA	1194
Tyr	His	Glu	Asn	Gly	Ser	Leu	Tyr	Asp	Phe	Leu	Lys	Cys	Ala	Thr	Leu	
				315					320					325		
GAC	ACC	AGA	GCC	CTA	CTC	AAG	TTA	GCT	TAT	TCT	GCT	GCT	TGT	GGT	CTG	1242
Asp	Thr	Arg	Ala	Leu	Leu	Lys	Leu	Ala	Tyr	Ser	Ala	Ala	Cys	Gly	Leu	
			330					335					340			

TGC CAC CTC CAC ACA GAA ATT TAT GGT ACC CAA GGG AAG CCT GCA ATT Cys His Leu His Thr Glu Ile Tyr Gly Thr Gln Gly Lys Pro Ala Ile 345 350 355	1290
GCT CAT CGA GAC CTG AAG AGC AAA AAC ATC CTT ATT AAG AAA AAT GGA Ala His Arg Asp Leu Lys Ser Lys Asn Ile Leu Ile Lys Lys Asn Gly 360 365 370	1338
AGT TGC TGT ATT GCT GAC CTG GGC CTA GCT GTT AAA TTC AAC AGT GAT Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala Val Lys Phe Asn Ser Asp 375 380 385 390	1386
ACA AAT GAA GTT GAC ATA CCC TTG AAT ACC AGG GTG GGC ACC AAG CGG Thr Asn Glu Val Asp Ile Pro Leu Asn Thr Arg Val Gly Thr Lys Arg 395 400 405	1434
TAC ATG GCT CCA GAA GTG CTG GAT GAA AGC CTG AAT AAA AAC CAT TTC Tyr Met Ala Pro Glu Val Leu Asp Glu Ser Leu Asn Lys Asn His Phe 410 415 420	1482
CAG CCC TAC ATC ATG GCT GAC ATC TAT AGC TTT GGT TTG ATC ATT TGG Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser Phe Gly Leu Ile Ile Trp 425 430 435	1530
GAA ATG GCT CGT CGT TGT ATT ACA GGA CGA ATC GTG GAG GAA TAT CAA Glu Met Ala Arg Arg Cys Ile Thr Gly Gly Ile Val Glu Glu Tyr Gln 440 445 450	1578
TTA CCA TAT TAC AAC ATG GTG CCC AGT GAC CCA TCC TAT GAG GAC ATG Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp Pro Ser Tyr Glu Asp Met 455 460 465 470	1626
CGT GAG GTT GTG TGT GTG AAA CGC TTG CGG CCA ATC GTG TCT AAC CGC Arg Glu Val Val Cys Val Lys Arg Leu Arg Pro Ile Val Ser Asn Arg 475 480 485	1674
TGG AAC AGC GAT GAA TGT CTT CGA GCA GTT TTG AAG CTA ATG TCA GAA Trp Asn Ser Asp Glu Cys Leu Arg Ala Val Leu Lys Leu Met Ser Glu 490 495 500	1722
TGT TGG GCC CAT AAT CCA GCC TCC AGA CTC ACA GCT TTG AGA ATC AAG Cys Trp Ala His Asn Pro Ala Ser Arg Leu Thr Ala Leu Arg Ile Lys 505 510 515	1770
AAG ACA CTT GCA AAA ATG GTT GAA TCC CAG GAT GTA AAG ATT Lys Thr Leu Ala Lys Met Val Glu Ser Gln Asp Val Lys Ile 520 525 530	1812
TGACAATTAA ACAATTTTGA GGGAGAATTT AGACTGCAAG AACCTTCTTCA CCCAAGGAAT	1872
GGGTGGGATT AGCATGGAAT AGGATGTTGA CTTGGTTTCC AGACTCCTTC CTCTACATCT	1932
TCACAGGCTG CTAACAGTAA ACCTTACCGT ACTCTACAGA ATACAAGATT GGAACCTTGA	1992
ACTTCAAACA TGTCATTCTT TATATATGAC AGCTTTGTTT TAATGTGGGG TTTTTTGT	2052
TGCTTTTTTT GTTTTGT	2070

## (2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 532 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

```

Met Thr Gln Leu Tyr Thr Tyr Ile Arg Leu Leu Gly Ala Cys Leu Phe
 1          5          10          15
Ile Ile Ser His Val Gln Gly Gln Asn Leu Asp Ser Met Leu His Gly
 20          25          30
Thr Gly Met Lys Ser Asp Leu Asp Gln Lys Lys Pro Glu Asn Gly Val
 35          40          45
Thr Leu Ala Pro Glu Asp Thr Leu Pro Phe Leu Lys Cys Tyr Cys Ser
 50          55          60
Gly His Cys Pro Asp Asp Ala Ile Asn Asn Thr Cys Ile Thr Asn Gly
 65          70          75          80
His Cys Phe Ala Ile Ile Glu Glu Asp Asp Gln Gly Glu Thr Thr Leu
 85          90          95
Thr Ser Gly Cys Met Lys Tyr Glu Gly Ser Asp Phe Gln Cys Lys Asp
100          105          110
Ser Pro Lys Ala Gln Leu Arg Arg Thr Ile Glu Cys Cys Arg Thr Asn
115          120          125
Leu Cys Asn Gln Tyr Leu Gln Pro Thr Leu Pro Pro Val Val Ile Gly
130          135          140
Pro Phe Phe Asp Gly Ser Ile Arg Trp Leu Val Val Leu Ile Ser Met
145          150          155          160
Ala Val Cys Ile Val Ala Met Ile Ile Phe Ser Ser Cys Phe Cys Tyr
165          170          175
Lys His Tyr Cys Lys Ser Ile Ser Ser Arg Gly Arg Tyr Asn Arg Asp
180          185          190
Leu Glu Gln Asp Glu Ala Phe Ile Pro Val Gly Glu Ser Leu Lys Asp
195          200          205
Leu Ile Asp Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu Pro Leu
210          215          220
Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Arg Gln Val
225          230          235          240
Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg Gly Glu
245          250          255

```

Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser Trp Phe  
 260 265 270  
 Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu Asn Ile  
 275 280 285  
 Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp Thr Gln  
 290 295 300  
 Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr Asp Phe  
 305 310 315 320  
 Leu Lys Cys Ala Thr Leu Asp Thr Arg Ala Leu Leu Lys Leu Ala Tyr  
 325 330 335  
 Ser Ala Ala Cys Gly Leu Cys His Leu His Thr Glu Ile Tyr Gly Thr  
 340 345 350  
 Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys Asn Ile  
 355 360 365  
 Leu Ile Lys Lys Asn Gly Ser Cys Cys Ile Ala Asp Leu Gly Leu Ala  
 370 375 380  
 Val Lys Phe Asn Ser Asp Thr Asn Glu Val Asp Ile Pro Leu Asn Thr  
 385 390 395 400  
 Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu Val Leu Asp Glu Ser  
 405 410 415  
 Leu Asn Lys Asn His Phe Gln Pro Tyr Ile Met Ala Asp Ile Tyr Ser  
 420 425 430  
~~Phe Gly Leu Ile Ile Trp Glu Met Ala Arg Arg Cys Ile Thr Gly Gly~~  
 435 440 445  
 Ile Val Glu Glu Tyr Gln Leu Pro Tyr Tyr Asn Met Val Pro Ser Asp  
 450 455 460  
 Pro Ser Tyr Glu Asp Met Arg Glu Val Val Cys Val Lys Arg Leu Arg  
 465 470 475 480  
 Pro Ile Val Ser Asn Arg Trp Asn Ser Asp Glu Cys Leu Arg Ala Val  
 485 490 495  
 Leu Lys Leu Met Ser Glu Cys Trp Ala His Asn Pro Ala Ser Arg Leu  
 500 505 510  
 Thr Ala Leu Arg Ile Lys Lys Thr Leu Ala Lys Met Val Glu Ser Gln  
 515 520 525  
 Asp Val Lys Ile  
 530

(2) INFORMATION FOR SEQ ID NO: 15:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2160 base pairs

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 10..1524

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

CGCGGTTAC ATG GCG GAG TCG GCC GGA GCC TCC TCC TTC TTC CCC CTT	48
Met Ala Glu Ser Ala Gly Ala Ser Ser Phe Phe Pro Leu	
1 5 10	
GTT GTC CTC CTG CTC GCC GGC AGC GGC GGG TCC GGG CCC CGG GGG ATC	96
Val Val Leu Leu Leu Ala Gly Ser Gly Gly Ser Gly Pro Arg Gly Ile	
15 20 25	
CAG GCT CTG CTG TGT GCG TGC ACC AGC TGC CTA CAG ACC AAC TAC ACC	144
Gln Ala Leu Leu Cys Ala Cys Thr Ser Cys Leu Gln Thr Asn Tyr Thr	
30 35 40 45	
TGT GAG ACA GAT GGG GCT TGC ATG GTC TCC ATC TTT AAC CTG GAT GGC	192
Cys Glu Thr Asp Gly Ala Cys Met Val Ser Ile Phe Asn Leu Asp Gly	
50 55 60	
GTG GAG CAC CAT GTA CGT ACC TGC ATC CCC AAG GTG GAG CTG GTT CCT	240
Val Glu His His Val Arg Thr Cys Ile Pro Lys Val Glu Leu Val Pro	
65 70 75	
GCT GGA AAG CCC TTC TAC TGC CTG AGT TCA GAG GAT CTG CGC AAC ACA	288
Ala Gly Lys Pro Phe Tyr Cys Leu Ser Ser Glu Asp Leu Arg Asn Thr	
80 85 90	
CAC TGC TGC TAT ATT GAC TTC TGC AAC AAG ATT GAC CTC AGG GTC CCC	336
His Cys Cys Tyr Ile Asp Phe Cys Asn Lys Ile Asp Leu Arg Val Pro	
95 100 105	
AGC GGA CAC CTC AAG GAG CCT GCG CAC CCC TCC ATG TGG GGC CCT GTG	384
Ser Gly His Leu Lys Glu Pro Ala His Pro Ser Met Trp Gly Pro Val	
110 115 120 125	
GAG CTG GTC GGC ATC ATC GCC GGC CCC GTC TTC CTC CTC TTC CTT ATC	432
Glu Leu Val Gly Ile Ile Ala Gly Pro Val Phe Leu Leu Phe Leu Ile	
130 135 140	





GCT CCT GAA GTC CTT GAC GAG ACA ATC AAC ATG AAG CAC TTT GAC TCC Ala Pro Glu Val Leu Asp Glu Thr Ile Asn Met Lys His Phe Asp Ser 385 390 395	1200
TTC AAA TGT GCC GAC ATC TAT GCC CTC GGG CTT GTC TAC TGG GAG ATT Phe Lys Cys Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp Glu Ile 400 405 410	1248
GCA CGA AGA TGC AAT TCT GGA GGA GTC CAT GAA GAC TAT CAA CTG CCG Ala Arg Arg Cys Asn Ser Gly Gly Val His Glu Asp Tyr Gln Leu Pro 415 420 425	1296
TAT TAC GAC TTA GTG CCC TCC GAC CCT TCC ATT GAG GAG ATG CGA AAG Tyr Tyr Asp Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met Arg Lys 430 435 440 445	1344
GTT GTA TGT GAC CAG AAG CTA CGG CCC AAT GTC CCC AAC TGG TGG CAG Val Val Cys Asp Gln Lys Leu Arg Pro Asn Val Pro Asn Trp Trp Gln 450 455 460	1392
AGT TAT GAG GCC TTG CGA GTG ATG GGA AAG ATG ATG CGG GAG TGC TGG Ser Tyr Glu Ala Leu Arg Val Met Gly Lys Met Met Arg Glu Cys Trp 465 470 475	1440
TAC GCC AAT GGT GCT GCC CGT CTG ACA GCT CTG CGC ATC AAG AAG ACT Tyr Ala Asn Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr 480 485 490	1488
CTG TCC CAG CTA AGC GTG CAG GAA GAT GTG AAG ATT TAAGCTGTTC Leu Ser Gln Leu Ser Val Gln Glu Asp Val Lys Ile 495 500 505	1534
CTCTGCCTAC ACAAAGAACC TGGGCAGTGA GGATGACTGC AGCCACCGTG CAAGCGTCGT	1594
GGAGGCCTAT CCTCTTGTTT CTGCCCCGCC CTCTGGCAGA GCCCTGGCCT GCAAGAGGGA	1654
CAGAGCCTGG GAGACGCGCG CACTCCCGTT GCGTTTGAGA CAGACACTTT TTATATTTAC	1714
CTCCTGATGG CATGGAGACC TGAGCAAATC ATGTAGTCAC TCAATGCCAC AACTCAAAC	1774
GCTTCAGTGG GAAGTACAGA GACCCAGTGC ATTGCGTGTG CAGGAGCGTG AGGTGCTGGG	1834
CTCGCCAGGA GCGGCCCCCA TACCTTGTGG TCCACTGGGC TGCAGGTTTT CCTCCAGGGA	1894
CCAGTCAACT GGCATCAAGA TATTGAGAGG AACCGGAAGT TTCTCCCTCC TTCCCGTAGC	1954
AGTCCTGAGC CACACCATCC TTCTCATGGA CATCCGGAGG ACTGCCCCTA GAGACACAAC	2014
CTGCTGCCTG TCTGTCCAGC CAAGTGCGCA TGTGCCGAGG TGTGTCCCAC ATTGTGCCTG	2074
GTCTGTGCCA CGCCCGTGTG TGTGTGTGTG TGTGTGAGTG AGTGTGTGTG TGTACACTTA	2134
ACCTGCTTGA GCTTCTGTGC ATGTGT	2160

## (2) INFORMATION FOR SEQ ID NO: 16:

- (1) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 505 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

```

Met  Ala  Glu  Ser  Ala  Gly  Ala  Ser  Ser  Phe  Phe  Pro  Leu  Val  Val  Leu
 1              5              10              15

Leu  Leu  Ala  Gly  Ser  Gly  Gly  Ser  Gly  Pro  Arg  Gly  Ile  Gln  Ala  Leu
              20              25              30

Leu  Cys  Ala  Cys  Thr  Ser  Cys  Leu  Gln  Thr  Asn  Tyr  Thr  Cys  Glu  Thr
              35              40              45

Asp  Gly  Ala  Cys  Met  Val  Ser  Ile  Phe  Asn  Leu  Asp  Gly  Val  Glu  His
      50              55              60

His  Val  Arg  Thr  Cys  Ile  Pro  Lys  Val  Glu  Leu  Val  Pro  Ala  Gly  Lys
 65              70              75              80

Pro  Phe  Tyr  Cys  Leu  Ser  Ser  Glu  Asp  Leu  Arg  Asn  Thr  His  Cys  Cys
              85              90              95

Tyr  Ile  Asp  Phe  Cys  Asn  Lys  Ile  Asp  Leu  Arg  Val  Pro  Ser  Gly  His
      100              105              110

Leu  Lys  Glu  Pro  Ala  His  Pro  Ser  Met  Trp  Gly  Pro  Val  Glu  Leu  Val
      115              120              125

Gly  Ile  Ile  Ala  Gly  Pro  Val  Phe  Leu  Leu  Phe  Leu  Ile  Ile  Ile  Ile
 130              135              140

Val  Phe  Leu  Val  Ile  Asn  Tyr  His  Gln  Arg  Val  Tyr  His  Asn  Arg  Gln
 145              150              155              160

Arg  Leu  Asp  Met  Glu  Asp  Pro  Ser  Cys  Glu  Met  Cys  Leu  Ser  Lys  Asp
      165              170              175

Lys  Thr  Leu  Gln  Asp  Leu  Val  Tyr  Asp  Leu  Ser  Thr  Ser  Gly  Ser  Gly
      180              185              190

Ser  Gly  Leu  Pro  Leu  Phe  Val  Gln  Arg  Thr  Val  Ala  Arg  Thr  Ile  Val
      195              200              205

Leu  Gln  Glu  Ile  Ile  Gly  Lys  Gly  Arg  Phe  Gly  Glu  Val  Trp  Arg  Gly
 210              215              220

Arg  Trp  Arg  Gly  Gly  Asp  Val  Ala  Val  Lys  Ile  Phe  Ser  Ser  Arg  Glu
 225              230              235              240

Glu  Arg  Ser  Trp  Phe  Arg  Glu  Ala  Glu  Ile  Tyr  Gln  Thr  Val  Met  Leu
      245              250              255

Arg  His  Glu  Asn  Ile  Leu  Gly  Phe  Ile  Ala  Ala  Asp  Asn  Lys  Asp  Asn
 260              265              270

```

74

Gly Thr Trp Thr Gln Leu Trp Leu Val Ser Asp Tyr His Glu His Gly  
 275 280 285  
 Ser Leu Phe Asp Tyr Leu Asn Arg Tyr Thr Val Thr Ile Glu Gly Met  
 290 295 300  
 Ile Lys Leu Ala Leu Ser Ala Ala Ser Gly Leu Ala His Leu His Met  
 305 310 315 320  
 Glu Ile Val Gly Thr Gln Gly Lys Pro Gly Ile Ala His Arg Asp Leu  
 325 330 335  
 Lys Ser Lys Asn Ile Leu Val Lys Lys Asn Gly Met Cys Ala Ile Ala  
 340 345 350  
 Asp Leu Gly Leu Ala Val Arg His Asp Ala Val Thr Asp Thr Ile Asp  
 355 360 365  
 Ile Ala Pro Asn Gln Arg Val Gly Thr Lys Arg Tyr Met Ala Pro Glu  
 370 375 380  
 Val Leu Asp Glu Thr Ile Asn Met Lys His Phe Asp Ser Phe Lys Cys  
 385 390 395 400  
 Ala Asp Ile Tyr Ala Leu Gly Leu Val Tyr Trp Glu Ile Ala Arg Arg  
 405 410 415  
 Cys Asn Ser Gly Gly Val His Glu Asp Tyr Gln Leu Pro Tyr Tyr Asp  
 420 425 430  
 Leu Val Pro Ser Asp Pro Ser Ile Glu Glu Met Arg Lys Val Val Cys  
 435 440 445  
 Asp Gln Lys Leu Arg Pro Asn Val Pro Asn Trp Trp Gln Ser Tyr Glu  
 450 455 460  
 Ala Leu Arg Val Met Gly Lys Met Met Arg Glu Cys Trp Tyr Ala Asn  
 465 470 475 480  
 Gly Ala Ala Arg Leu Thr Ala Leu Arg Ile Lys Lys Thr Leu Ser Gln  
 485 490 495  
 Leu Ser Val Gln Glu Asp Val Lys Ile  
 500 505

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1952 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: unknown
  - (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 187..1692

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

AAGCGGCGGC AGAAGTTGCC GCGGTGGTGC TCGTAGTGAG GCGCGGGAGG ACCCGGGACC	60
TGGGAAGCGG CGGCGGGTTA ACTTCGGCTG AATCACAACC ATTTGGCGCT GAGCTATGAC	120
AAGAGAGCAA ACAAAAAGTT AAAGGAGCAA CCGGGCCATA AGTGAAGAGA GAAGTTTATT	180
GATAAC ATG CTC TTA CGA AGC TCT GGA AAA TTA AAT GTG GGC ACC AAG	228
Met Leu Leu Arg Ser Ser Gly Lys Leu Asn Val Gly Thr Lys	
1 5 10	
AAG GAG GAT GGA GAG AGT ACA GCC CCC ACC CCT CGG CCC AAG ATC CTA	276
Lys Glu Asp Gly Glu Ser Thr Ala Pro Thr Pro Arg Pro Lys Ile Leu	
15 20 25 30	
CGT TGT AAA TGC CAC CAC CAC TGT CCG GAA GAC TCA GTC AAC AAT ATC	324
Arg Cys Lys Cys His His His Cys Pro Glu Asp Ser Val Asn Asn Ile	
35 40 45	
TGC AGC ACA GAT GGG TAC TGC TTC ACG ATG ATA GAA GAA GAT GAC TCT	372
Cys Ser Thr Asp Gly Tyr Cys Phe Thr Met Ile Glu Glu Asp Asp Ser	
50 55 60	
GGA ATG CCT GTT GTC ACC TCT GGA TGT CTA GGA CTA GAA GGG TCA GAT	420
Gly Met Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp	
65 70 75	
TTT CAA TGT CGT GAC ACT CCC ATT CCT CAT CAA AGA AGA TCA ATT GAA	468
Phe Gln Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu	
80 85 90	
TGC TGC ACA GAA AGG AAT GAG TGT AAT AAA GAC CTC CAC CCC ACT CTG	516
Cys Cys Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu	
95 100 105 110	
CCT CCT CTC AAG GAC AGA GAT TTT GTT GAT GGG CCC ATA CAC CAC AAG	564
Pro Pro Leu Lys Asp Arg Asp Phe Val Asp Gly Pro Ile His His Lys	
115 120 125	
GCC TTG CTT ATC TCT GTG ACT GTC TGT AGT TTA CTC TTG GTC CTC ATT	612
Ala Leu Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile	
130 135 140	
ATT TTA TTC TGT TAC TTC AGG TAT AAA AGA CAA GAA GCC CGA CCT CGG	660
Ile Leu Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Ala Arg Pro Arg	
145 150 155	

TAC Tyr 160	AGC Ser 160	ATT Ile 160	GGG Gly 160	CTG Leu 160	GAG Glu 165	CAG Gln 165	GAC Asp 165	GAG Glu 165	ACA Thr 165	TAC Tyr 170	ATT Ile 170	CCT Pro 170	CCT Pro 170	GGA Gly 170	GAG Glu 170	708
TCC Ser 175	CTG Leu 175	AGA Arg 175	GAC Asp 175	TTG Leu 180	ATC Ile 180	GAG Glu 180	CAG Gln 180	TCT Ser 185	CAG Gln 185	AGC Ser 185	TCG Ser 185	GGA Gly 190	AGT Ser 190	GGA Gly 190	TCA Ser 190	756
GGC Gly 195	CTC Leu 195	CCT Pro 195	CTG Leu 195	CTG Leu 195	GTC Val 195	CAA Gln 200	AGG Arg 200	ACA Thr 200	ATA Ile 200	GCT Ala 200	AAG Lys 205	CAA Gln 205	ATT Ile 205	CAG Gln 205	ATG Met 205	804
GTG Val 210	AAG Lys 210	CAG Gln 210	ATT Ile 210	GGA Gly 210	AAA Lys 215	GGC Gly 215	CGC Arg 215	TAT Tyr 215	GGC Gly 215	GAG Glu 215	GTG Val 220	TGG Trp 220	ATG Met 220	GGA Gly 220	AAG Lys 220	852
TGG Trp 225	CGT Arg 225	GGA Gly 225	GAA Glu 225	AAG Lys 230	GTG Val 230	GCT Ala 230	GTG Val 230	AAA Lys 235	GTG Val 235	TTC Phe 235	TTC Phe 235	ACC Thr 235	ACG Thr 235	GAG Glu 235	GAA Glu 235	900
GCC Ala 240	AGC Ser 240	TGG Trp 240	TTC Phe 240	CGA Arg 245	GAG Glu 245	ACT Thr 245	GAG Glu 245	ATA Ile 250	TAT Tyr 250	CAG Gln 250	ACG Thr 250	GTC Val 250	CTG Leu 250	ATG Met 250	CGG Arg 250	948
CAT His 255	GAG Glu 255	AAT Asn 255	ATT Ile 255	CTG Leu 260	GGG Gly 260	TTC Phe 260	ATT Ile 260	GCT Ala 265	GCA Ala 265	GAT Asp 265	ATC Ile 265	AAA Lys 270	GGG Gly 270	ACT Thr 270	GGG Gly 270	996
TCC Ser 275	TGG Trp 275	ACT Thr 275	CAG Gln 275	TTG Leu 275	TAC Tyr 275	CTC Leu 280	ATC Ile 280	ACA Thr 280	GAC Asp 280	TAT Tyr 285	CAT His 285	GAA Glu 285	AAC Asn 285	GGC Gly 285	TCC Ser 285	1044
CTT Leu 290	TAT Tyr 290	GAC Asp 290	TAT Tyr 290	CTG Leu 290	AAA Lys 295	TCC Ser 295	ACC Thr 295	ACC Thr 295	TTA Leu 295	GAC Asp 295	GCA Ala 300	AAG Lys 300	TCC Ser 300	ATG Met 300	CTG Leu 300	1092
AAG Lys 305	CTA Leu 305	GCC Ala 305	TAC Tyr 305	TCC Ser 305	TCT Ser 310	GTC Val 310	AGC Ser 310	CGC Gly 310	CTA Leu 315	TGC Cys 315	CAT His 315	TTA Leu 315	CAC His 315	ACG Thr 315	GAA Glu 315	1140
ATC Ile 320	TTT Phe 320	AGC Ser 320	ACT Thr 320	CAA Gln 325	GGC Gly 325	AAG Lys 325	CCA Pro 325	GCA Ala 330	ATC Ile 330	GCC Ala 330	CAT His 330	CGA Arg 330	GAC Asp 330	TTG Leu 330	AAA Lys 330	1188
AGT Ser 335	AAA Lys 335	AAC Asn 335	ATC Ile 340	CTG Leu 340	GTG Val 340	AAG Lys 340	AAA Lys 345	AAT Asn 345	GGA Gly 345	ACT Thr 345	TGC Cys 345	TGC Cys 345	ATA Ile 350	GCA Ala 350	GAC Asp 350	1236
CTG Leu 355	GGC Gly 355	TTG Leu 355	GCT Ala 355	GTC Val 355	AAG Lys 355	TTC Phe 360	ATT Ile 360	AGT Ser 360	GAC Asp 360	ACA Thr 365	AAT Asn 365	GAG Glu 365	GTT Val 365	GAC Asp 365	ATC Ile 365	1284
CCA Pro 370	CCC Pro 370	AAC Asn 370	ACC Thr 370	CGG Arg 375	GTT Val 375	GGC Gly 375	ACC Thr 375	AAG Lys 375	CGC Arg 375	TAT Tyr 380	ATG Met 380	CCT Pro 380	CCA Pro 380	GAA Glu 380	GTG Val 380	1332
CTG Leu 385	GAC Asp 385	GAG Glu 385	AGC Ser 385	TTG Leu 390	AAT Asn 390	AGA Arg 390	AAC Asn 390	CAT His 390	TTC Phe 390	CAG Gln 395	TCC Ser 395	TAC Tyr 395	ATT Ile 395	ATG Met 395	GCT Ala 395	1380

GAC ATG TAC AGC TTT GGA CTC ATC CTC TGG GAG ATT GCA AGG AGA TGT Asp Met Tyr Ser Phe Gly Leu Ile Leu Trp Glu Ile Ala Arg Arg Cys 400 405 410	1428
GTT TCT GGA GGT ATA GTG GAA GAA TAC CAG CTT CCC TAT CAC GAC CTG Val Ser Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu 415 420 425 430	1476
GTG CCC AGT GAC CCT TCT TAT GAG GAC ATG AGA GAA ATT GTG TGC ATG Val Pro Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Met 435 440 445	1524
AAG AAG TTA CGG CCT TCA TTC CCC AAT CGA TGG AGC AGT GAT GAG TGT Lys Lys Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys 450 455 460	1572
CTC AGG CAG ATG GGG AAG CTT ATG ACA GAG TGC TGG GCG CAG AAT CCT Leu Arg Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala Gln Asn Pro 465 470 475	1620
GCC TCC AGG CTG ACG GCC CTG AGA GTT AAG AAA ACC CTT GCC AAA ATG Ala Ser Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met 480 485 490	1668
TCA GAG TCC CAG GAC ATT AAA CTC TGACGTCAGA TACTTGTGGA CAGAGCAAGA Ser Glu Ser Gln Asp Ile Lys Leu 495 500	1722
ATTTACACAGA AGCATCGTTA GCCCAAGCCT TGAACGTTAG CCTACTGCCC AGTGAGTTCA	1782
GACTTTCCTG GAAGAGAGCA CCGTGGGCAG ACACAGAGGA ACCCAGAAAC ACGGATTCAT	1842
CATGGCTTTC TGAGGAGGAG AAAGTGTGTTG GGTAACTTGT TCAAGATATG ATGCATGTTG	1902
CTTTCTAAGA AAGCCCTGTA TTTTGAATTA CCATTTTTTTT ATAAAAAAA	1952

## (2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 502 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Met	Leu	Leu	Arg	Ser	Ser	Gly	Lys	Leu	Asn	Val	Gly	Thr	Lys	Lys	Glu	
1				5					10					15		
Asp	Gly	Glu	Ser	Thr	Ala	Pro	Thr	Pro	Arg	Pro	Lys	Ile	Leu	Arg	Cys	
			20					25					30			
Lys	Cys	His	His	His	Cys	Pro	Glu	Asp	Ser	Val	Asn	Asn	Ile	Cys	Ser	
		35					40					45				
Thr	Asp	Gly	Tyr	Cys	Phe	Thr	Met	Ile	Glu	Glu	Asp	Asp	Ser	Gly	Met	
	50						55					60				

78

Pro Val Val Thr Ser Gly Cys Leu Gly Leu Glu Gly Ser Asp Phe Gln  
 65 70 75 80  
 Cys Arg Asp Thr Pro Ile Pro His Gln Arg Arg Ser Ile Glu Cys Cys  
 85 90 95  
 Thr Glu Arg Asn Glu Cys Asn Lys Asp Leu His Pro Thr Leu Pro Pro  
 100 105 110  
 Leu Lys Asp Arg Asp Phe Val Asp Gly Pro Ile His His Lys Ala Leu  
 115 120 125  
 Leu Ile Ser Val Thr Val Cys Ser Leu Leu Leu Val Leu Ile Ile Leu  
 130 135 140  
 Phe Cys Tyr Phe Arg Tyr Lys Arg Gln Glu Ala Arg Pro Arg Tyr Ser  
 145 150 155 160  
 Ile Gly Leu Glu Gln Asp Glu Thr Tyr Ile Pro Pro Gly Glu Ser Leu  
 165 170 175  
 Arg Asp Leu Ile Glu Gln Ser Gln Ser Ser Gly Ser Gly Ser Gly Leu  
 180 185 190  
 Pro Leu Leu Val Gln Arg Thr Ile Ala Lys Gln Ile Gln Met Val Lys  
 195 200 205  
 Gln Ile Gly Lys Gly Arg Tyr Gly Glu Val Trp Met Gly Lys Trp Arg  
 210 215 220  
 Gly Glu Lys Val Ala Val Lys Val Phe Phe Thr Thr Glu Glu Ala Ser  
 225 230 235 240  
 Trp Phe Arg Glu Thr Glu Ile Tyr Gln Thr Val Leu Met Arg His Glu  
 245 250 255  
 Asn Ile Leu Gly Phe Ile Ala Ala Asp Ile Lys Gly Thr Gly Ser Trp  
 260 265 270  
 Thr Gln Leu Tyr Leu Ile Thr Asp Tyr His Glu Asn Gly Ser Leu Tyr  
 275 280 285  
 Asp Tyr Leu Lys Ser Thr Thr Leu Asp Ala Lys Ser Met Leu Lys Leu  
 290 295 300  
 Ala Tyr Ser Ser Val Ser Gly Leu Cys His Leu His Thr Glu Ile Phe  
 305 310 315 320  
 Ser Thr Gln Gly Lys Pro Ala Ile Ala His Arg Asp Leu Lys Ser Lys  
 325 330 335  
 Asn Ile Leu Val Lys Lys Asn Gly Thr Cys Cys Ile Ala Asp Leu Gly  
 340 345 350  
 Leu Ala Val Lys Phe Ile Ser Asp Thr Asn Glu Val Asp Ile Pro Pro  
 355 360 365  
 Asn Thr Arg Val Gly Thr Lys Arg Tyr Met Pro Pro Glu Val Leu Asp  
 370 375 380



79

Glu Ser Leu Asn Arg Asn His Phe Gln Ser Tyr Ile Met Ala Asp Met  
 385 390 395 400  
 Tyr Ser Phe Gly Leu Ile Leu Trp Glu Ile Ala Arg Arg Cys Val Ser  
 405 410 415  
 Gly Gly Ile Val Glu Glu Tyr Gln Leu Pro Tyr His Asp Leu Val Pro  
 420 425 430  
 Ser Asp Pro Ser Tyr Glu Asp Met Arg Glu Ile Val Cys Met Lys Lys  
 435 440 445  
 Leu Arg Pro Ser Phe Pro Asn Arg Trp Ser Ser Asp Glu Cys Leu Arg  
 450 455 460  
 Gln Met Gly Lys Leu Met Thr Glu Cys Trp Ala Gln Asn Pro Ala Ser  
 465 470 475 480  
 Arg Leu Thr Ala Leu Arg Val Lys Lys Thr Leu Ala Lys Met Ser Glu  
 485 490 495  
 Ser Gln Asp Ile Lys Leu  
 500

## (2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

GCGGATCCTG TTGTGAAGGN AATATGTC

28

## (2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GGGATCCGTC GCAGTCAAAA TTTT

24

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 26 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

GCGGATCCGC GATATATTAA AAGCAA

26

(2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

CGGAATTCTG GTGCCATATA

20

(2) INFORMATION FOR SEQ ID NO: 23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

ATTCAAGGGC ACATCAACTT CATTGTGTC ACTGTTG

37

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GCGGATCCAC CATGGCGGAG TCGGCC

26

(2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

AACACCGGGC CGGCGATGAT

20

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

Gly Xaa Gly Xaa Xaa Gly  
1 5

(2) INFORMATION FOR SEQ ID NO: 27:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

Asp Phe Lys Ser Arg Asn  
1 5

(2) INFORMATION FOR SEQ ID NO: 28:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

Asp Leu Lys Ser Lys Asn  
1 5

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

Gly Thr Lys Arg Tyr Met  
1 5

CLAIMS

1. An isolated protein having a serine/threonine kinase domain, a DFKSRN or DLKSKN sequence in subdomain VIB and/or a GTKRYM sequence in subdomain VIII.
- 5 2. A protein according to claim 1, which additionally comprises an ATP-binding sequence that is Gly-Xaa-Gly-Xaa-Xaa-Gly in subdomain I, and a Lys residue in subdomain II.
3. An isolated protein having a serine/threonine kinase domain which has more than 50% identity to the kinase  
10 domain of any of the amino-acid sequences identified herein as SEQ ID. Nos. 2, 4, 6, 8, 10, 12, 14, 16 and 18.
4. A protein according to claim 3, wherein the identity is more than 60%.
5. A protein according to any preceding claim, having  
15 serine/threonine kinase activity.
6. An isolated protein having all or part of any of the amino-acid sequences identified herein as SEQ. ID Nos. 2, 4, 6, 8, 10, 12, 14, 16 and 18, and activin receptor type I functionality.
- 20 7. An isolated protein having an amino-acid sequence corresponding to part or all of the amino-acid sequence of an activin type I receptor, and wherein the protein has at least one of the following characteristics:-
  - (i) serine/threonine kinase activity;
  - 25 (ii) activin-binding activity; and
  - (iii) activin type II receptor interaction.
8. An isolated protein having all or part of any of the amino-acid sequences identified herein as SEQ. ID Nos. 2, 4, 6, 8, 10, 12, 14, 16 and 18, and TGF- $\beta$ -type I receptor  
30 functionality.
9. An isolated protein having an amino-acid sequence corresponding to part or all of the amino-acid sequence of a TGF- $\beta$ -type I receptor, and wherein the protein has at least one of the following characteristics:
  - 35 (i) serine/threonine kinase activity;
  - (ii) TGF- $\beta$ -binding activity; and
  - (iii) TGF- $\beta$ -type II receptor interaction.

10. A protein according to any of claims 1 to 5, having all or part of the amino-acid sequence identified herein as SEQ ID No. 2.
11. A protein according to any of claims 1 to 7, having  
5 all or part of the amino-acid sequence identified herein as SEQ ID No. 4.
12. A protein according to any of claims 1 to 5, having serine/threonine kinase activity and all or part of the amino-acid sequence identified herein as SEQ ID No. 6.
- 10 13. A protein according to any of claims 1 to 7, having all or part of the amino-acid sequence identified herein as SEQ ID No. 8.
14. A protein according to any of claims 1 to 5, 8 and 9, having all or part of the amino-acid sequence identified  
15 herein as SEQ ID No. 10.
15. A protein according to any of claims 1 to 5, having all or part of the amino-acid sequence identified herein as SEQ ID No. 12.
16. A protein according to any of claims 1 to 5, having  
20 all or part of the amino-acid sequence identified herein as SEQ ID No. 14.
17. A protein according to any of claims 1 to 7, having all or part of the amino-acid sequence identified herein as SEQ ID No. 16.
- 25 18. A protein according to any of claims 1 to 5, having all or part of the amino-acid sequence identified herein as SEQ ID No. 18.
19. A protein according to any preceding claim, that is a soluble receptor.
- 30 20. An antibody which binds specifically to a protein as defined in any of claims 1 to 19 and not to at least one other such protein.
21. An isolated nucleic acid molecule which codes for, or is complementary to a nucleic acid molecule which codes  
35 for, a protein as defined in any of claims 1 to 19.
22. A recombinant nucleic acid molecule comprising at least two heterologous sequences, one of which codes for,

or is complementary to a nucleic acid molecule which codes for, a protein as defined in any of claims 1 to 19.

23. A molecule according to claim 21 or claim 22, wherein the protein is a TGF- $\beta$ -type I receptor.

5 24. A molecule according to claim 21 or claim 22, wherein the protein is an activin receptor.

25. A DNA or RNA/mRNA molecule according to any of claims 21 to 24.

10 26. A molecule according to any of claims 20 to 24, which additionally comprises, operably associated with the coding sequence, a sequence adapted to allow expression of the protein.

27. A host comprising a molecule according to claim 26, which is capable of expressing the protein.

15 28. A host according to claim 27, which comprises PAE cells.

29. A host according to claim 27 or claim 28, transfected with the Chim A receptor plasmid.

20 30. A product according to any preceding claim, for therapeutic or diagnostic use.

31. Use of a product according to any of claims 1 to 29, for the manufacture of a medicament for use in treating a condition associated with TGF activity.

1/11

cons.aa	<u>G G</u>	<u>G V</u>	<u>A K</u>	<u>E</u>
hTGfBR-II	LDTLVGKGRFAEVYKAKLKQNTSEQFETVAVKIFPYDHYASWKORKDIFSDINLKHENILQF			
mActR-IIB	LLEIKARGRFGCVWKAQLMN-----DFVAVKIKPLQDKQSWQSEREIFSTPGMTHENLLQF			
mActR-II	LLEVKARGRFGCVWKAQLLN-----EYVAVKIFPIQDKQSWQNEYEVYSIPGMTHENILQF			
daf-1	LTVRVGSGRFGNVSRGDYRG-----EAVAVKVFNAIDEPAPFHKEIEIFETRMLRHPNVLR			
subdomains	I		II	III IV

hTGfBR-II	LTAEERKTEL GKQYWLITAFHAKGNLQEYLTRHVISWEDLRNVGSSSLARGLSHLSDHTP-C
mActR-IIB	IAAEKRGSNLEVELWLITAFHDKGSLIDYLKGNITWNECHVAETMSRGISYLVHEDVPWCR
mActR-II	IGAERGTSDVDLWLITAFHEKGSLSDFLKANVVSWNELCHIAETMARGLAYLHEDIPGLK
daf-1	IGSDRVDTGFTLWLVIEYHPSGSLHDFLENTVNIETYYNLMRSTASGLAFLHNQIGGSK
subdomains	V VI-A

cons.aa	<u>DLK</u>	<u>N</u>	<u>DFG</u>
hTGfBR-II	-GRPKMPIVHRDLKSSNIVKNDLTCCLCDFGLSLRL--		-GPYSSVDDLANSQGQVGTARYMAP
mActR-IIB	GECHKPSIAHRDFKSKNVLLKSDLTAVLADFGGLAVRF--		-EPGKPPGD--THGQVGTRRYMAP
mActR-II	-DGHKPAISHRDIKSKNVLLKNNLTACIADFGGLALKF--		-EAGKSAGD--THGQVGTRRYMAP
daf-1	-ESNKPAMAHRDIKSKNIMYKNDLTCAIGDLGLSLSKPEDAASDIAN--		-ENYKCGTVRYLAP
subdomains	VI-B		VII VIII

Fig. 1



2/11

---

a.a            C C E G N M C  
5' GCGGATCCTGTTGTGAAGGNAATATGTG 3' Fig. 2A  
BAMHI C C G C

---

a.a            V A V K I F  
5' GCGGATCCGTCGCAGTCAAATTTT 3' Fig. 2B  
BamHI G C G G C  
T T T A

---

a.a            R D I K S K N  
5' GCGGATCCGCGATATTAAAAGCAA 3' Fig. 2C  
BAMHI A C C GTCT  
G A

---

a.a            E P A M Y  
5' CGGAATTCTGGTGCCATATA Fig. 2D  
EcoRI G G G  
A A





**Fig. 3 contd.**

6/11

K	M	N	L	T	A	C	I	A	D	F	G	L	A	L	K	F	E	A	G	K	S	A	G	D	D	L	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	E	G	Actr-II	
K	S	D	L	T	A	V	L	A	D	F	G	L	A	V	R	L	F	E	P	G	K	P	P	G	D	D	L	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	E	G	Actr-IIIB
K	M	D	L	T	C	C	L	C	D	F	G	L	S	L	R	L	D	S	A	T	S	V	D	D	L	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	E	S	Tbr-II		
K	K	N	G	T	C	C	I	A	D	L	C	G	L	A	V	R	H	D	S	A	T	D	T	I	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	Tbr-I/ALK-S		
K	K	N	L	Q	C	C	I	A	D	L	C	G	L	A	V	M	H	S	Q	S	S	D	Y	L	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	ALK-1		
K	K	N	G	S	C	C	I	A	D	L	C	G	L	A	V	M	H	S	Q	S	S	D	Y	L	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	ALK-2		
K	K	N	G	S	C	C	I	A	D	L	C	G	L	A	V	K	F	N	S	S	D	T	N	E	V	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	ALK-3	
K	K	N	G	M	C	A	I	A	D	L	G	L	A	V	R	H	D	A	V	T	D	T	N	E	V	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	ALK-4	
K	K	N	G	T	C	C	I	A	D	L	G	L	A	V	K	F	I	S	D	T	N	E	V	D	I	A	P	H	S	T	H	G	Q	V	G	T	R	R	Y	M	A	P	E	V	L	D	ALK-6			

VIII

VII


A	I	M	F	Q	R	-	D	A	F	L	R	I	D	M	Y	A	M	G	L	V	L	W	E	L	A	S	R	C	T	A	A	D	G	P	V	D	E	Y	M	L	P	F	E	E	Actr-II
A	I	M	F	Q	R	-	D	A	F	L	R	I	D	M	Y	A	M	G	L	V	L	W	E	L	A	S	R	C	T	A	A	D	G	P	V	D	E	Y	M	L	P	F	E	E	Actr-IIIB
R	M	L	E	N	A	E	S	F	K	Q	I	D	I	D	Y	S	M	A	L	V	L	W	E	M	T	S	R	C	N	A	V	-	G	E	V	K	D	Y	E	P	F	G	S	Tbr-II	
S	I	N	K	H	F	E	S	F	K	R	A	D	I	I	Y	A	M	G	L	V	F	W	E	I	A	R	R	C	S	I	-	G	I	H	E	D	Y	Q	L	P	Y	D	Tbr-I/ALK-S		
Q	I	R	T	D	C	F	E	S	Y	K	W	T	D	I	M	A	F	G	L	V	L	W	E	I	A	R	R	T	I	V	-	N	G	I	V	E	D	Y	R	P	F	Y	D	ALK-1	
T	I	Q	V	D	C	F	D	S	Y	K	R	V	D	I	M	A	F	G	L	V	L	W	E	I	A	R	R	M	V	S	-	N	G	I	V	E	D	Y	R	P	F	Y	D	ALK-2	
S	L	N	K	N	H	F	Q	P	Y	I	M	A	D	I	Y	S	E	G	L	I	I	W	E	M	A	R	R	C	I	T	-	G	I	V	E	E	Y	Q	L	P	Y	N	ALK-3		
T	I	N	K	N	H	F	D	S	F	K	C	A	D	I	Y	A	L	G	L	V	Y	W	E	I	A	R	R	C	N	S	-	G	G	V	M	E	E	Y	Q	L	P	Y	D	ALK-4	
S	L	N	R	N	H	E	Q	S	Y	I	M	A	D	M	Y	S	E	G	L	I	L	W	E	I	A	R	R	C	V	S	-	G	G	I	V	E	E	Y	Q	L	P	Y	H	D	ALK-6

IX

X

E	I	G	Q	H	P	S	L	E	D	M	Q	E	V	V	V	H	K	K	K	R	P	V	L	R	D	Y	W	Q	K	H	A	G	M	A	M	L	C	E	T	I	E	E	C	W	Actr-II					
E	I	G	Q	H	P	S	L	E	E	L	Q	E	V	V	V	L	R	D	H	W	L	K	T	I	K	D	H	W	L	K	H	A	G	M	A	M	L	C	V	T	I	E	E	C	W	Actr-IIIB				
K	V	R	E	M	P	C	V	E	S	M	K	D	N	V	V	C	E	I	P	S	F	W	I	P	S	F	W	Q	K	N	Q	I	Q	M	V	C	E	T	L	I	E	E	C	W	Tsr-II					
L	V	P	S	D	P	S	V	E	E	M	R	K	V	V	V	C	E	Q	K	L	R	P	N	I	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Tsr-I/ALK-S
V	V	P	N	D	P	S	F	E	D	M	K	K	V	V	V	C	V	D	Q	Q	T	P	N	I	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Alk-1
M	V	P	N	D	P	S	F	E	D	M	R	K	V	V	V	C	V	D	Q	Q	R	P	N	I	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Alk-2
L	V	P	S	D	P	S	F	E	D	M	R	K	V	V	V	C	V	D	Q	Q	R	P	N	I	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Alk-3
L	V	P	S	D	P	S	F	E	D	M	R	K	V	V	V	C	V	D	Q	Q	R	P	N	I	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Alk-4
L	V	P	S	D	P	S	F	E	D	M	R	E	I	V	C	M	K	K	L	R	P	S	F	P	N	R	W	L	A	D	P	V	L	S	G	L	R	V	M	A	K	I	M	M	R	E	C	W	Alk-6	

Fig. 3 contd.



XI

P K E S S L (S13) A C T R - I I  
 P K E S S I (S36) A C T R - I I B  
 K (S67) T 6 R - I I

Fig. 3 contd.

8/11




Fig. 4





10/11

ALK-2	ALK-3	ALK-4	ALK-5	ActR-II	ActR-IIB	TβR-II	daf-1	
79	60	61	63	40	40	37	39	ALK-1
	63	64	65	41	39	37	39	ALK-2
		63	65	41	38	37	39	ALK-3
			90	41	40	39	42	ALK-4
				42	40	41	43	ALK-5
					78	48	35	ActR-II
						47	32	ActR-IIB
							34	TβR-II

Fig. 6




Fig. 7

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**